

**Anti-BiP/GRP78 Antibody**  
**Mouse Anti Human Monoclonal Antibody**  
**Catalog # AP53389****Specification**

---

**Anti-BiP/GRP78 Antibody - Product Information**

Application	IP, WB, ICC
Primary Accession	<a href="#">P11021</a>
Other Accession	<a href="#">NM_005347</a>
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Immunogen	Purified recombinant human BiP/GRP78 protein expressed in E.coli.
Purification	Affinity purified
Calculated MW	78 KDa

**Anti-BiP/GRP78 Antibody - Additional Information****Gene ID** 3309**Other Names**

78 kDa glucose regulated protein;78 kDa glucose-regulated protein;AL022860;AU019543;BiP;D2Wsu141e;D2Wsu17e;Endoplasmic reticulum lumenal Ca(2+)-binding protein grp78;Endoplasmic reticulum lumenal Ca2+ binding protein grp78;FLJ26106;Glucose Regulated Protein 78kDa;GRP 78;GRP-78;GRP78;GRP78\_HUMAN;Heat shock 70 kDa protein 5;Heat Shock 70kDa Protein 5;Hsce70;HSPA 5;HSPA5;Immunoglobulin Heavy Chain Binding Protein;Immunoglobulin heavy chain-binding protein;mBiP;MIF2;Sez7.

**Dilution**

WB~~1:1000

**Format**

PBS(pH 7.4) containing with 0.09% (W/V) sodium azide and 50% glycerol.

**Storage**

Store at -20 °C.Stable for 12 months from date of receipt

**Anti-BiP/GRP78 Antibody - Protein Information****Name** HSPA5 ([HGNC:5238](#))**Function**

Endoplasmic reticulum chaperone that plays a key role in protein folding and quality control in the endoplasmic reticulum lumen (PubMed:<a href="http://www.uniprot.org/citations/2294010" target="\_blank">2294010</a>, PubMed:<a href="http://www.uniprot.org/citations/23769672" target="\_blank">23769672</a>, PubMed:<a href="http://www.uniprot.org/citations/23990668"

target="\_blank">23990668</a>, PubMed:<a href="http://www.uniprot.org/citations/28332555" target="\_blank">28332555</a>). Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10/ERdj5, probably to facilitate the release of DNAJC10/ERdj5 from its substrate (By similarity). Acts as a key repressor of the ERN1/IRE1-mediated unfolded protein response (UPR) (PubMed:<a href="http://www.uniprot.org/citations/1550958" target="\_blank">1550958</a>, PubMed:<a href="http://www.uniprot.org/citations/19538957" target="\_blank">19538957</a>). In the unstressed endoplasmic reticulum, recruited by DNAJB9/ERdj4 to the luminal region of ERN1/IRE1, leading to disrupt the dimerization of ERN1/IRE1, thereby inactivating ERN1/IRE1 (By similarity). Accumulation of misfolded protein in the endoplasmic reticulum causes release of HSPA5/BiP from ERN1/IRE1, allowing homodimerization and subsequent activation of ERN1/IRE1 (By similarity). Plays an auxiliary role in post-translational transport of small presecretory proteins across endoplasmic reticulum (ER). May function as an allosteric modulator for SEC61 channel-forming translocon complex, likely cooperating with SEC62 to enable the productive insertion of these precursors into SEC61 channel. Appears to specifically regulate translocation of precursors having inhibitory residues in their mature region that weaken channel gating. May also play a role in apoptosis and cell proliferation (PubMed:<a href="http://www.uniprot.org/citations/26045166" target="\_blank">26045166</a>).

#### Cellular Location


Endoplasmic reticulum lumen. Melanosome. Cytoplasm {ECO:0000250|UniProtKB:P20029}. Cell surface Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:12643545). Localizes to the cell surface of epithelial cells in response to high levels of free iron (PubMed:20484814, PubMed:24355926, PubMed:27159390)


#### Anti-BiP/GRP78 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-BiP/GRP78 Antibody - Images






Immunoprecipitation of BiP/GRP78 from HeLa cell extracts using BiP/GRP78 Mouse mAb. Western blot was performed using BiP/GRP78 Mouse mAb.



Western blot analysis of extracts from 3T3, A549, C6 and HeLa cell lysates using BiP/GRP78 mouse mAb (1:1000 diluted). Predicted band size: 72KDa. Observed band size: 72KDa.



Immunofluorescent analysis of HeLa cells fixed by anhydrous methanol at -20°C and using BiP/GRP78 mouse mAb (dilution 1:200).

**Anti-BiP/GRP78 Antibody - Background**

Probably plays a role in facilitating the assembly of multimeric protein complexes inside the endoplasmic reticulum. Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10, probably to facilitate