

Anti-SIRT6 Antibody
Mouse Anti Human Monoclonal Antibody
Catalog # AP53392

Specification

Anti-SIRT6 Antibody - Product Information

Application	IP, WB, ICC
Primary Accession	Q8N6T7
Other Accession	NM_016539
Reactivity	Human, Mouse, Rat, Monkey
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Immunogen	Purified recombinant human SIRT6 protein expressed in E.coli.
Purification	Affinity purified
Calculated MW	42 KDa

Anti-SIRT6 Antibody - Additional Information

Gene ID 51548

Other Names

2810449N18Rik;AI043036;Mono ADP ribosyltransferase sirtuin 6;NAD-dependent protein deacetylase sirtuin-6;Regulatory protein SIR2 homolog 6;Regulatory protein SIR2 homolog;SIR2 like 6;SIR2 like protein 6;Sir2 related protein type 6;SIR2-like protein 6;SIR2

Dilution

WB~~1:500
ICC~~1:100

Format

PBS(pH 7.4) containing with 0.09% (W/V) sodium azide and 50% glycerol.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

Anti-SIRT6 Antibody - Protein Information

Name SIRT6 {ECO:0000303|PubMed:10873683, ECO:0000312|HGNC:HGNC:14934}

Function

NAD-dependent protein deacetylase, deacylase and mono-ADP- ribosyltransferase that plays an essential role in DNA damage repair, telomere maintenance, metabolic homeostasis, inflammation, tumorigenesis and aging (PubMed:18337721, PubMed:19135889, PubMed:19625767, PubMed:19625767).

[21680843](http://www.uniprot.org/citations/21680843), PubMed: [23217706](http://www.uniprot.org/citations/23217706), PubMed: [23653361](http://www.uniprot.org/citations/23653361), PubMed: [24052263](http://www.uniprot.org/citations/24052263), PubMed: [27322069](http://www.uniprot.org/citations/27322069), PubMed: [27180906](http://www.uniprot.org/citations/27180906), PubMed: [21362626](http://www.uniprot.org/citations/21362626), PubMed: [23552949](http://www.uniprot.org/citations/23552949), PubMed: [30374165](http://www.uniprot.org/citations/30374165), PubMed: [29555651](http://www.uniprot.org/citations/29555651)). Displays protein- lysine deacetylase or defatty-acylase (demyrystoylase and depalmitoylase) activity, depending on the context (PubMed: [24052263](http://www.uniprot.org/citations/24052263) target="_blank">24052263, PubMed: [27322069](http://www.uniprot.org/citations/27322069) target="_blank">27322069, PubMed: [23552949](http://www.uniprot.org/citations/23552949) target="_blank">23552949). Acts as a key histone deacetylase by catalyzing deacetylation of histone H3 at 'Lys-9', 'Lys-18' and 'Lys- 56' (H3K9ac, H3K18ac and H3K56ac, respectively), suppressing target gene expression of several transcription factors, including NF-kappa-B (PubMed: [19625767](http://www.uniprot.org/citations/19625767) target="_blank">19625767, PubMed: [24012758](http://www.uniprot.org/citations/24012758) target="_blank">24012758, PubMed: [23892288](http://www.uniprot.org/citations/23892288) target="_blank">23892288, PubMed: [23911928](http://www.uniprot.org/citations/23911928) target="_blank">23911928, PubMed: [27043296](http://www.uniprot.org/citations/27043296) target="_blank">27043296, PubMed: [26898756](http://www.uniprot.org/citations/26898756) target="_blank">26898756, PubMed: [27180906](http://www.uniprot.org/citations/27180906) target="_blank">27180906, PubMed: [33067423](http://www.uniprot.org/citations/33067423) target="_blank">33067423, PubMed: [21362626](http://www.uniprot.org/citations/21362626) target="_blank">21362626, PubMed: [30374165](http://www.uniprot.org/citations/30374165) target="_blank">30374165, PubMed: [26456828](http://www.uniprot.org/citations/26456828) target="_blank">26456828). Acts as an inhibitor of transcription elongation by mediating deacetylation of H3K9ac and H3K56ac, preventing release of NELFE from chromatin and causing transcriptional pausing (By similarity). Involved in DNA repair by promoting double-strand break (DSB) repair: acts as a DSB sensor by recognizing and binding DSB sites, leading to (1) recruitment of DNA repair proteins, such as SMARCA5/SNF2H, and (2) deacetylation of histone H3K9ac and H3K56ac (PubMed: [23911928](http://www.uniprot.org/citations/23911928) target="_blank">23911928, PubMed: [31995034](http://www.uniprot.org/citations/31995034) target="_blank">31995034, PubMed: [32538779](http://www.uniprot.org/citations/32538779) target="_blank">32538779). SIRT6 participation to DSB repair is probably involved in extension of life span (By similarity). Also promotes DNA repair by deacetylating non-histone proteins, such as DDB2 and p53/TP53 (PubMed: [32789493](http://www.uniprot.org/citations/32789493) target="_blank">32789493, PubMed: [29474172](http://www.uniprot.org/citations/29474172) target="_blank">29474172). Specifically deacetylates H3K18ac at pericentric heterochromatin, thereby maintaining pericentric heterochromatin silencing at centromeres and protecting against genomic instability and cellular senescence (PubMed: [27043296](http://www.uniprot.org/citations/27043296) target="_blank">27043296). Involved in telomere maintenance by catalyzing deacetylation of histone H3 in telomeric chromatin, regulating telomere position effect and telomere movement in response to DNA damage (PubMed: [18337721](http://www.uniprot.org/citations/18337721) target="_blank">18337721, PubMed: [19625767](http://www.uniprot.org/citations/19625767) target="_blank">19625767, PubMed: [21847107](http://www.uniprot.org/citations/21847107) target="_blank">21847107). Required for embryonic stem cell differentiation by mediating histone deacetylation of H3K9ac (PubMed: [25915124](http://www.uniprot.org/citations/25915124) target="_blank">25915124, PubMed: [29555651](http://www.uniprot.org/citations/29555651) target="_blank">29555651). Plays a major role in metabolism by regulating processes such as glycolysis, gluconeogenesis, insulin secretion and lipid metabolism (PubMed: [24012758](http://www.uniprot.org/citations/24012758) target="_blank">24012758, PubMed: [26787900](http://www.uniprot.org/citations/26787900) target="_blank">26787900). Inhibits glycolysis via histone deacetylase activity and by acting as a corepressor of the transcription factor HIF1A, thereby controlling the expression of multiple glycolytic genes (By similarity). Has

tumor suppressor activity by repressing glycolysis, thereby inhibiting the Warburg effect (PubMed:23217706). Also regulates glycolysis and tumorigenesis by mediating deacetylation and nuclear export of non-histone proteins, such as isoform M2 of PKM (PKM2) (PubMed:26787900). Acts as a negative regulator of gluconeogenesis by mediating deacetylation of non-histone proteins, such as FOXO1 and KAT2A/GCN5 (PubMed:23142079, PubMed:25009184). Promotes beta-oxidation of fatty acids during fasting by catalyzing deacetylation of NCOA2, inducing coactivation of PPARA (By similarity). Acts as a regulator of lipid catabolism in brown adipocytes, both by catalyzing deacetylation of histones and non-histone proteins, such as FOXO1 (By similarity). Also acts as a regulator of circadian rhythms, both by regulating expression of clock-controlled genes involved in lipid and carbohydrate metabolism, and by catalyzing deacetylation of PER2 (By similarity). The defatty-acylase activity is specifically involved in regulation of protein secretion (PubMed:24052263, PubMed:23552949, PubMed:27322069, PubMed:28406396). Has high activity toward long-chain fatty acyl groups and mediates protein-lysine demyristoylation and depalmitoylation of target proteins, such as RRAS2 and TNF, thereby regulating their secretion (PubMed:23552949, PubMed:28406396). Also acts as a mono-ADP- ribosyltransferase by mediating mono-ADP-ribosylation of PARP1, TRIM28/KAP1 or SMARCC2/BAF170 (PubMed:21680843, PubMed:22753495, PubMed:27568560, PubMed:27322069). Mono-ADP-ribosyltransferase activity is involved in DNA repair, cellular senescence, repression of LINE-1 retrotransposon elements and regulation of transcription (PubMed:21680843, PubMed:22753495, PubMed:27568560).

Cellular Location


Nucleus. Chromosome. Chromosome, telomere. Endoplasmic reticulum. Note=Predominantly nuclear (PubMed:18337721). Associated with pericentric heterochromatin and telomeric heterochromatin regions (PubMed:18337721, PubMed:27043296) Localizes to DNA damage sites: directly recognizes and binds double-strand breaks (DSBs) sites via a tunnel-like structure that has high affinity for DSBs (PubMed:21680843, PubMed:23911928, PubMed:27568560, PubMed:31995034, PubMed:32538779). A fraction localizes to the endoplasmic reticulum (PubMed:23552949).

Anti-SIRT6 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-SIRT6 Antibody - Images



Immunoprecipitation analysis of HeLa cell lysates using SIRT6 mouse mAb.



Western blot analysis of extracts from Jurkat, MCF7, HCT116, K562, COS7, C2C12, 3T3 and HeLa cell lysates using SIRT6 mouse mAb (1:500 diluted). Predicted band size: 42,36KDa. Observed band size: 42,36KDa.



Immunofluorescent analysis of Hela cells fixed by 4% paraformaldehyde and using SIRT6 mouse mAb (dilution 1:100).

Anti-SIRT6 Antibody - Background

NAD-dependent protein deacetylase. Has deacetylase activity towards histone H3K9Ac and H3K56Ac. Modulates acetylation of histone H3 in telomeric chromatin during the S-phase of the cell cycle. Deacetylates histone H3K9Ac at NF-kappa-B target promoters and