

Anti-Toll-like Receptor 3 Antibody
Mouse Monoclonal Antibody
Catalog # AP53439**Specification**

Anti-Toll-like Receptor 3 Antibody - Product Information

Application	WB
Primary Accession	O15455
Other Accession	NM_003265
Reactivity	Transfected
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Immunogen	Purified recombinant human Toll-Like Receptor 3 protein fragments expressed in E.coli.
Purification	Affinity purified

Anti-Toll-like Receptor 3 Antibody - Additional Information**Gene ID** 7098**Other Names**

CD283 ; CD283 antigen ; IIAE2 ; TLR 3 ; Tlr3 ; TLR3_HUMAN ; Toll Like Receptor 3 ; Toll-like receptor 3.

Dilution

WB~~1:1000

Format

Purified mouse monoclonal antibody in PBS(pH 7.4) containing with 0.09% (W/V) sodium azide and 50% glycerol.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

Anti-Toll-like Receptor 3 Antibody - Protein Information**Name** TLR3 ([HGNC:11849](#))**Function**

Key component of innate and adaptive immunity. TLRs (Toll- like receptors) control host immune response against pathogens through recognition of molecular patterns specific to microorganisms. TLR3 is a nucleotide-sensing TLR which is activated by double-stranded RNA, a sign of viral infection. Acts via the adapter TRIF/TICAM1, leading to NF-kappa-B activation, IRF3 nuclear translocation, cytokine secretion and the inflammatory response.

Cellular Location

Endoplasmic reticulum membrane; Single-pass type I membrane protein. Endosome membrane. Early endosome

Tissue Location


Expressed at high level in placenta and pancreas. Also detected in CD11c+ immature dendritic cells. Only expressed in dendritic cells and not in other leukocytes, including monocyte precursors. TLR3 is the TLR that is expressed most strongly in the brain, especially in astrocytes, glia, and neurons

Anti-Toll-like Receptor 3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Toll-like Receptor 3 Antibody - Images



Western blot detection of Toll-like Receptor 3 in CHO-K1 cell lysate [B] and CHO-K1 transfected by Toll-like Receptor 3 [A] cell lysate using Toll-like Receptor 3 mouse mAb (1:1000 diluted). Predicted band size: 50KDa. Observed band size: 50KDa.

Anti-Toll-like Receptor 3 Antibody - Background

Key component of innate and adaptive immunity. TLRs (Toll-like receptors) control host immune response against pathogens through recognition of molecular patterns specific of microorganisms. TLR3 is a nucleotide-sensing TLR which is activated by double-str