

DYNC1LI2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5478c

Specification

DYNC1LI2 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	O43237
Other Accession	O4R5P6 , NP_006132.1
Reactivity	Human
Predicted	Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	54099
Antigen Region	152-180

DYNC1LI2 Antibody (Center) - Additional Information

Gene ID 1783

Other Names

Cytoplasmic dynein 1 light intermediate chain 2, Dynein light intermediate chain 2, cytosolic, LIC-2, LIC53/55, DYNC1LI2, DNCL12, LIC2

Target/Specificity

This DYNC1LI2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 152-180 amino acids from the Central region of human DYNC1LI2.

Dilution

WB~~1:2000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DYNC1LI2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

DYNC1LI2 Antibody (Center) - Protein Information

Name DYNC1LI2 ([HGNC:2966](#))

Synonyms DNCL12, LIC2

Function Acts as one of several non-catalytic accessory components of the cytoplasmic dynein 1 complex that are thought to be involved in linking dynein to cargos and to adapter proteins that regulate dynein function. Cytoplasmic dynein 1 acts as a motor for the intracellular retrograde motility of vesicles and organelles along microtubules. May play a role in binding dynein to membranous organelles or chromosomes.


Cellular Location

Cytoplasm, cytoskeleton


DYNC1LI2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DYNC1LI2 Antibody (Center) - Images

All lanes : Anti-DYNC1LI2 Antibody (Center) at 1:2000 dilution Lane 1: Human brain lysate Lane 2: U-87 MG whole cell lysate Lane 3: SK-OV-3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 54 kDa Blocking/Dilution buffer: 5% NFD/MTBST.



DYNC1LI2 Antibody (Center) (Cat. #AP5478c) immunohistochemistry analysis in formalin fixed and paraffin embedded human cervix carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the DYNC1LI2 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

DYNC1LI2 Antibody (Center) - Background

Cytoplasmic dynein is a microtubule-associated motor protein (Hughes et al., 1995 [PubMed 7738094]). See DYNC1H1 (MIM 600112) for general information about dyneins.

DYNC1LI2 Antibody (Center) - References

Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)
Olsen, J.V., et al. Cell 127(3):635-648(2006)
Beausoleil, S.A., et al. Nat. Biotechnol. 24(10):1285-1292(2006)