

## **RGS22 Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP5631c

## **Specification**

## **RGS22 Antibody (Center) - Product Information**

WB, IHC-P, FC,E Application **Primary Accession O8NE09** Other Accession NP 056483.3 Human, Mouse Reactivity Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Antigen Region 388-417

# **RGS22 Antibody (Center) - Additional Information**

#### **Gene ID 26166**

#### **Other Names**

Regulator of G-protein signaling 22, RGS22, RGS22

# Target/Specificity

This RGS22 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 388-417 amino acids from the Central region of human RGS22.

#### **Dilution**

WB~~1:1000 IHC-P~~1:50~100 FC~~1:10~50

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

## **Precautions**

RGS22 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## **RGS22 Antibody (Center) - Protein Information**

### Name RGS22

Function Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits



thereby driving them into their inactive GDP-bound form.

#### **Cellular Location**

Cytoplasm. Nucleus. Note=Expressed in the cytoplasm of spermatogonia and spermatocytes. In spermatids, also expressed in the nucleus

#### **Tissue Location**


Testis-specific. Expressed in Leydig cells and spermatogenic cells from the spermatogonia to spermatid stages (at protein level).

# **RGS22 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## **RGS22 Antibody (Center) - Images**




RGS22 Antibody (Center) (Cat. #AP5631c) western blot analysis in mouse heart tissue lysates (15ug/lane). This demonstrates the RGS22 antibody detected the RGS22 protein (arrow).





RGS22 Antibody (Center) (Cat. #AP5631c) immunohistochemistry analysis in formalin fixed and paraffin embedded human testis tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the RGS22 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



RGS22 Antibody (Center) (Cat. #AP5631c) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

# **RGS22 Antibody (Center) - Background**

RGS22 inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form (By similarity).

# **RGS22 Antibody (Center) - References**

Hu, Y., et al. Biol. Reprod. 79(6):1021-1029(2008)