

MURC Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5646a

Specification

MURC Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q5BKX8
Other Accession	NP_001018126.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	87-116

MURC Antibody (N-term) - Additional Information

Gene ID 347273

Other Names

Muscle-related coiled-coil protein, Muscle-restricted coiled-coil protein, MURC

Target/Specificity

This MURC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 87-116 amino acids from the N-terminal region of human MURC.

Dilution

WB~~1:1000
IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MURC Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MURC Antibody (N-term) - Protein Information

Name CAVIN4 ([HGNC:33742](#))

Synonyms MURC

Function Modulates the morphology of formed caveolae in cardiomyocytes, but is not required for caveolar formation. Facilitates the recruitment of MAPK1/3 to caveolae within cardiomyocytes and regulates alpha-1 adrenergic receptor-induced hypertrophic responses in cardiomyocytes through MAPK1/3 activation. Contributes to proper membrane localization and stabilization of caveolin-3 (CAV3) in cardiomyocytes (By similarity). Induces RHOA activation and activates NPPA transcription and myofibrillar organization through the Rho/ROCK signaling pathway (PubMed:[18332105](#)).

Cellular Location


Cytoplasm, myofibril, sarcomere {ECO:0000250|UniProtKB:A2AMM0}. Cytoplasm {ECO:0000250|UniProtKB:A2AMM0}. Cytoplasm, cytosol {ECO:0000250|UniProtKB:A2AMM0}. Cell membrane, sarcolemma {ECO:0000250|UniProtKB:A2AMM0}. Membrane, caveola. Cell membrane. Note=In cardiomyocytes, accumulates in the Z-line of the sarcomere. In vascular smooth muscle cells, detected diffusely throughout the cytoplasm. Localizes in the caveolae in a caveolin-dependent manner. {ECO:0000250|UniProtKB:A2AMM0}

MURC Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MURC Antibody (N-term) - Images



Anti-MURC Antibody (N-term) at 1:1000 dilution + human heart lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 42 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



MURC Antibody (N-term) (Cat. #AP5646a) immunohistochemistry analysis in formalin fixed and paraffin embedded human lung carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the MURC Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

MURC Antibody (N-term) - Background

MURC induces RHOA activation and activates NPPA transcription and myofibrillar organization through the Rho/ROCK signaling pathway.

MURC Antibody (N-term) - References

Tagawa, M., et al. Am. J. Physiol., Cell Physiol. 295 (2), C490-C498 (2008) :
Ogata, T., et al. Mol. Cell. Biol. 28(10):3424-3436(2008)