

NDST2 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5759b

Specification

NDST2 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P52849
Other Accession	NP_003626.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	817-844

NDST2 Antibody (C-term) - Additional Information

Gene ID 8509

Other Names

Bifunctional heparan sulfate N-deacetylase/N-sulfotransferase 2, Glucosaminyl N-deacetylase/N-sulfotransferase 2, NDST-2, N-heparan sulfate sulfotransferase 2, N-HSST 2, Heparan sulfate N-deacetylase 2, 3---, Heparan sulfate N-sulfotransferase 2, 282-, NDST2, HSST2

Target/Specificity

This NDST2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 817-844 amino acids of human NDST2.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NDST2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NDST2 Antibody (C-term) - Protein Information

Name NDST2

Synonyms HSST2

Function Essential bifunctional enzyme that catalyzes both the N- deacetylation and the N-sulfation of glucosamine (GlcNAc) of the glycosaminoglycan in heparan sulfate. Modifies the GlcNAc-GlcA disaccharide repeating sugar backbone to make N-sulfated heparosan, a prerequisite substrate for later modifications in heparin biosynthesis. Plays a role in determining the extent and pattern of sulfation of heparan sulfate. Required for the exosomal release of SDCBP, CD63 and syndecan (PubMed:[22660413](#)).

Cellular Location


Golgi apparatus membrane; Single- pass type II membrane protein

NDST2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NDST2 Antibody (C-term) - Images



NDST2 Antibody (C-term) (Cat. #AP5759b) western blot analysis in mouse cerebellum tissue lysates (15ug/lane). This demonstrates the NDST2 antibody detected NDST2 protein (arrow).

NDST2 Antibody (C-term) - Background

NDST2 is a member of the N-deacetylase/N-sulfotransferase subfamily of the sulfotransferase 1 proteins. The encoded enzyme has dual functions in processing glucosamine and heparin polymers, including N-deacetylation and N-sulfation. The encoded protein may be localized to the Golgi.

NDST2 Antibody (C-term) - References

Duncan, M.B., et al. Biochem. Biophys. Res. Commun. 339(4):1232-1237(2006)
Grupe, A., et al. Am. J. Hum. Genet. 78(1):78-88(2006)

Carter, N.M., et al. J. Cell. Sci. 116 (PT 17), 3591-3600 (2003)

NDST2 Antibody (C-term) - Citations

- [Modulation of heparan sulfate biosynthesis by sodium butyrate in recombinant CHO cells.](#)
- [Bioengineering murine mastocytoma cells to produce anticoagulant heparin.](#)
- [Bioengineered Chinese hamster ovary cells with Golgi-targeted 3-O-sulfotransferase-1 biosynthesize heparan sulfate with an antithrombin-binding site.](#)