

FAM26E Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5814B**Specification**

FAM26E Antibody (C-term) - Product Information

Application	WB, FC,E
Primary Accession	Q8N5C1
Other Accession	NP_714922.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35170
Antigen Region	214-243

FAM26E Antibody (C-term) - Additional Information**Gene ID** 254228**Other Names**

Protein FAM26E, FAM26E, C6orf188

Target/Specificity

This FAM26E antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 214-243 amino acids from the C-terminal region of human FAM26E.

Dilution

WB~~1:1000

FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FAM26E Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

FAM26E Antibody (C-term) - Protein Information**Name** CALHM5 ([HGNC:21568](#))**Synonyms** C6orf188, FAM26E

Function Pore-forming subunit of a voltage-gated ion channel.

Cellular Location


Membrane; Multi-pass membrane protein

FAM26E Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FAM26E Antibody (C-term) - Images



FAM26E Antibody (C-term) (Cat. #AP5814b) western blot analysis in mouse stomach tissue lysates (15ug/lane). This demonstrates the FAM26E antibody detected FAM26E protein (arrow).



FAM26E Antibody (C-term) (Cat. #AP5814b) flow cytometric analysis of 293 cells (right)

histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

FAM26E Antibody (C-term) - References

Mungall, A.J., et al. Nature 425(6960):805-811(2003)