

## VKORC1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP5859A

## **Specification**

## VKORC1 Antibody (N-term) - Product Information

Application WB, IHC-P,E
Primary Accession O9BOB6

Other Accession <u>Q6B4I2</u>, <u>NP 076869.1</u>

Reactivity
Predicted
Host
Clonality
Isotype
Antigen Region

Human
Bovine
Rabbit
Polyclonal
Rabbit IgG
15-41

## VKORC1 Antibody (N-term) - Additional Information

#### **Gene ID 79001**

### **Other Names**

Vitamin K epoxide reductase complex subunit 1, Vitamin K1 2, 3-epoxide reductase subunit 1, VKORC1, VKOR

## Target/Specificity

This VKORC1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 15-41 amino acids from the N-terminal region of human VKORC1.

## **Dilution**

WB~~1:1000 IHC-P~~1:10~50

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

VKORC1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

# VKORC1 Antibody (N-term) - Protein Information

Name VKORC1 {ECO:0000303|PubMed:14765194, ECO:0000312|HGNC:HGNC:23663}





**Function** Involved in vitamin K metabolism. Catalytic subunit of the vitamin K epoxide reductase (VKOR) complex which reduces inactive vitamin K 2,3-epoxide to active vitamin K. Vitamin K is required for the gamma-carboxylation of various proteins, including clotting factors, and is required for normal blood coagulation, but also for normal bone development.

#### **Cellular Location**

Endoplasmic reticulum membrane; Multi-pass membrane protein

### **Tissue Location**


Expressed at highest levels in fetal and adult liver, followed by fetal heart, kidney, and lung, adult heart, and pancreas.

# VKORC1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# VKORC1 Antibody (N-term) - Images



Anti-VKORC1 Antibody (N-term) at 1:1000 dilution + Human lung whole tissue lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution. Predicted band size : 18 kDa Blocking/Dilution buffer: 5% NFDM/TBST.





VKORC1 Antibody (N-term) (Cat. #AP5859a) immunohistochemistry analysis in formalin fixed and paraffin embedded human hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the VKORC1 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.