

**SIAH2 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP5878c****Specification**

---

**SIAH2 Antibody (Center) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">O43255</a>
Other Accession	<a href="#">Q9I8X5</a> , <a href="#">Q8R4T2</a> , <a href="#">Q06986</a> , <a href="#">Q7SYL3</a> , <a href="#">NP_005058.3</a>
Reactivity	Human
Predicted	Zebrafish, Mouse, Rat, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	216-244

**SIAH2 Antibody (Center) - Additional Information****Gene ID** 6478**Other Names**

E3 ubiquitin-protein ligase SIAH2, 632-, Seven in absentia homolog 2, Siah-2, hSiah2, SIAH2

**Target/Specificity**

This SIAH2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 216-244 amino acids from the Central region of human SIAH2.

**Dilution**WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

SIAH2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**SIAH2 Antibody (Center) - Protein Information****Name** SIAH2

**Function** E3 ubiquitin-protein ligase that mediates ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:[9334332](#), PubMed:[11483518](#), PubMed:[19224863](#)). E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates (PubMed:[9334332](#), PubMed:[11483518](#), PubMed:[19224863](#)). Mediates E3 ubiquitin ligase activity either through direct binding to substrates or by functioning as the essential RING domain subunit of larger E3 complexes (PubMed:[9334332](#), PubMed:[11483518](#), PubMed:[19224863](#)). Triggers the ubiquitin-mediated degradation of many substrates, including proteins involved in transcription regulation (GPS2, POU2AF1, PML, NCOR1), a cell surface receptor (DCC), an antiapoptotic protein (BAG1), and a protein involved in synaptic vesicle function in neurons (SYP) (PubMed:[9334332](#), PubMed:[11483518](#), PubMed:[19224863](#)). Mediates ubiquitination and proteasomal degradation of DYRK2 in response to hypoxia (PubMed:[22878263](#)). It is thereby involved in apoptosis, tumor suppression, cell cycle, transcription and signaling processes (PubMed:[9334332](#), PubMed:[11483518](#), PubMed:[19224863](#), PubMed:[22878263](#)). Has some overlapping function with SIAH1 (PubMed:[9334332](#), PubMed:[11483518](#), PubMed:[19224863](#)). Triggers the ubiquitin-mediated degradation of TRAF2, whereas SIAH1 does not (PubMed:[12411493](#)). Promotes monoubiquitination of SNCA (PubMed:[19224863](#)). Regulates cellular clock function via ubiquitination of the circadian transcriptional repressors NR1D1 and NR1D2 leading to their proteasomal degradation (PubMed:[26392558](#)). Plays an important role in mediating the rhythmic degradation/clearance of NR1D1 and NR1D2 contributing to their circadian profile of protein abundance (PubMed:[26392558](#)). Mediates ubiquitination and degradation of EGLN2 and EGLN3 in response to the unfolded protein response (UPR), leading to their degradation and subsequent stabilization of ATF4 (By similarity). Also part of the Wnt signaling pathway in which it mediates the Wnt-induced ubiquitin-mediated proteasomal degradation of AXIN1.

**Cellular Location**

Cytoplasm. Nucleus Note=Predominantly cytoplasmic. Partially nuclear

**Tissue Location**


Widely expressed at low level.

**SIAH2 Antibody (Center) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**SIAH2 Antibody (Center) - Images**



SIAH2 Antibody (Center) (Cat. #AP5878c) western blot analysis in T47D cell line lysates (35ug/lane). This demonstrates the SIAH2 antibody detected the SIAH2 protein (arrow).



SIAH2 Antibody (Center) (Cat. #AP5878c) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the SIAH2 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



SIAH2 Antibody (Center) (Cat. #AP5878c) flow cytometric analysis of T47D cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.