

M Sirt3 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6242B

Specification

M Sirt3 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q8R104
Other Accession	NP_071878.2
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	304-334

M Sirt3 Antibody (C-term) - Additional Information

Gene ID 64384

Other Names

NAD-dependent protein deacetylase sirtuin-3, 351-, Regulatory protein SIR2 homolog 3, SIR2-like protein 3, mSIR2L3, Sirt3, Sir2l3

Target/Specificity

This Sirt3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 304-334 amino acids from the C-terminal region of Mouse Sirt3.

Dilution

WB~~1:2000
IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

M Sirt3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

M Sirt3 Antibody (C-term) - Protein Information

Name Sirt3

Synonyms Sir2l3

Function NAD-dependent protein deacetylase (PubMed:[23835326](#), PubMed:[17923681](#), PubMed:[18794531](#), PubMed:[21172655](#), PubMed:[26620563](#)). Activates or deactivates mitochondrial target proteins by deacetylating key lysine residues (PubMed:[23835326](#), PubMed:[17923681](#), PubMed:[18794531](#), PubMed:[21172655](#)). Known targets include ACSS1, IDH, GDH, PDHA1, SOD2, LCAD, SDHA and the ATP synthase subunit ATP5PO (PubMed:[16790548](#), PubMed:[18794531](#), PubMed:[21172655](#)). Contributes to the regulation of the cellular energy metabolism (PubMed:[23835326](#), PubMed:[36804859](#)). Important for regulating tissue-specific ATP levels (PubMed:[18794531](#), PubMed:[24252090](#)). In response to metabolic stress, deacetylates transcription factor FOXO3 and recruits FOXO3 and mitochondrial RNA polymerase POLRMT to mtDNA to promote mtDNA transcription (PubMed:[23283301](#)). Acts as a regulator of ceramide metabolism by mediating deacetylation of ceramide synthases CERS1, CERS2 and CERS6, thereby increasing their activity and promoting mitochondrial ceramide accumulation (PubMed:[26620563](#)). Regulates hepatic lipogenesis. Uses NAD(+) substrate imported by SLC25A47, triggering downstream activation of PRKAA1/AMPK-alpha signaling cascade that ultimately downregulates sterol regulatory element-binding protein (SREBP) transcriptional activities and ATP-consuming lipogenesis to restore cellular energy balance.

Cellular Location

[Isoform L]: Mitochondrion matrix

Tissue Location


Expressed in cardiomyocytes (at protein level) (PubMed:11056054, PubMed:35959657). Expressed in the brain, liver, kidney and testes (PubMed:11056054). Expressed in skeletal muscles (at protein level) (PubMed:23283301, PubMed:23835326)


M Sirt3 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

M Sirt3 Antibody (C-term) - Images







Anti-Mouse Sirt3 Antibody (C-term) at 1:2000 dilution + mouse liver lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Anti-Sirt3 Antibody (C-term) at 1:64000 dilution + mouse liver lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-Sirt3 Antibody (C-term) at 1:2000 dilution + mouse liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-Sirt3 Antibody (C-term) at 1:2000 dilution + CHO whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Sirt3 Antibody (C-term) (Cat. #AP6242b) IHC analysis in formalin fixed and paraffin embedded

human breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the Sirt3 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

M Sirt3 Antibody (C-term) - Background

SIRT3 is a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The SIRT3 is included in class I of the sirtuin family.

M Sirt3 Antibody (C-term) - References

Hirschey, M.D., et al. Nature 464(7285):121-125(2010)

Pillai, V.B., et al. J. Biol. Chem. 285(5):3133-3144(2010)

Kim, H.S., et al. Cancer Cell 17(1):41-52(2010)

M Sirt3 Antibody (C-term) - Citations

- [Mouse SIRT3 attenuates hypertrophy-related lipid accumulation in the heart through the deacetylation of LCAD.](#)
- [Sirt3 protects in vitro-fertilized mouse preimplantation embryos against oxidative stress-induced p53-mediated developmental arrest.](#)