

VEGF Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6290b**Specification**

VEGF Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P15692
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	43597

VEGF Antibody (C-term) - Additional Information**Gene ID** 7422**Other Names**

Vascular endothelial growth factor A, VEGF-A, Vascular permeability factor, VPF, VEGFA, VEGF

Target/Specificity

This VEGF antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the C-terminal region of human VEGF Antibody.

Dilution

WB~~1:2000

IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

VEGF Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

VEGF Antibody (C-term) - Protein Information**Name** VEGFA**Synonyms** VEGF**Function** [N-VEGF]: Participates in the induction of key genes involved in the response to hypoxia

and in the induction of angiogenesis such as HIF1A (PubMed:[35455969](#)). Involved in protecting cells from hypoxia- mediated cell death (By similarity).

Cellular Location

[N-VEGF]: Cytoplasm. Nucleus. Note=Cytoplasmic in normoxic conditions and localizes to the nucleus under hypoxic conditions [Isoform L-VEGF189]: Endoplasmic reticulum. Golgi apparatus. Secreted, extracellular space, extracellular matrix [Isoform VEGF165]: Secreted

Tissue Location


Higher expression in pituitary tumors than the pituitary gland. [Isoform VEGF165]: Widely expressed. [Isoform VEGF206]: Not widely expressed.

VEGF Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


VEGF Antibody (C-term) - Images



All lanes : Anti-VEGF Antibody (C-term) at 1:2000 dilution Lane 1: MCF-7 whole cell lysate Lane 2: PC-3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 27 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-VEGF Antibody (C-term) at 1:2000 dilution Lane 1: MCF-7 whole cell lysate Lane 2: PC-3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 27 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



Formalin-fixed and paraffin-embedded human breast carcinoma tissue reacted with VEGF Antibody (C-term)(Cat.#AP6290b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

VEGF Antibody (C-term) - Background

Embryonic vascular system undergoes a series of complex, highly regulated series of events involving differentiation, migration and association of primitive endothelial cells. This process is termed vasculogenesis. A further remodeling of the primitive vascular system forms the mature cardiovascular system. This process is known as angiogenesis (sprouting of new capillary vessels from pre-existing vasculature). Angiogenesis accounts for the formation of vasculature into previously avascular organs such as brain and kidney. Angiogenic activity in the adult is required during the normal tissue repair, and for the remodeling of the female reproductive organs (ovulation and placental development). Certain pathological conditions, such as tumor growth and diabetic retinopathy, also require angiogenesis. Study of tumor angiogenesis has led to the identification of several proteins including basic fibroblast growth factor (bFGF) and vascular endothelial growth factor. VEGF acts by interacting with a family of largely endothelial-specific receptor tyrosine kinases that includes VEGFR-1 (flt-1), VEGFR-2 (flk-1/KDR), and VEGFR-3/Flt-4. Disruption of VEGFRs interferes with differentiation of endothelial cells and it is lethal for the embryo.

VEGF is a heparin-binding glycoprotein that is secreted as a homodimer of 45 kDa.

VEGF Antibody (C-term) - Citations

- [A novel LC-MS/MS approach to the pharmacokinetic study of free and bound aflibercept simultaneously](#)
- [Inhibition of ATM reverses EMT and decreases metastatic potential of cisplatin-resistant lung cancer cells through JAK/STAT3/PD-L1 pathway.](#)