

**FLI1 Antibody (N-term K67)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP6514D****Specification**

---

**FLI1 Antibody (N-term K67) - Product Information**

|                   |                                                 |
|-------------------|-------------------------------------------------|
| Application       | WB, IHC-P,E                                     |
| Primary Accession | <a href="#">Q01543</a>                          |
| Other Accession   | <a href="#">P26323</a> , <a href="#">Q29RS8</a> |
| Reactivity        | Human                                           |
| Predicted         | Bovine, Mouse                                   |
| Host              | Rabbit                                          |
| Clonality         | Polyclonal                                      |
| Isotype           | Rabbit IgG                                      |
| Calculated MW     | 50982                                           |
| Antigen Region    | 52-80                                           |

**FLI1 Antibody (N-term K67) - Additional Information****Gene ID** 2313**Other Names**

Friend leukemia integration 1 transcription factor, Proto-oncogene Fli-1, Transcription factor ERGB, FLI1

**Target/Specificity**

This FLI1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 52-80 amino acids from the N-terminal region of human FLI1.

**Dilution**WB~~1:1000  
IHC-P~~1:50~100**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

FLI1 Antibody (N-term K67) is for research use only and not for use in diagnostic or therapeutic procedures.

**FLI1 Antibody (N-term K67) - Protein Information****Name** FLI1

**Function** Sequence-specific transcriptional activator (PubMed:[24100448](#), PubMed:[26316623](#), PubMed:[28255014](#)). Recognizes the DNA sequence 5'- C[CA]GGAAGT-3'.

**Cellular Location**


Nucleus.

**FLI1 Antibody (N-term K67) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**FLI1 Antibody (N-term K67) - Images**



Western blot analysis of FLI1 antibody (N-term) (Cat. #AP6514d) in 293 cell line lysates (35ug/lane). FLI1 (arrow) was detected using the purified Pab.



FLI-1 Antibody (N-term) (Cat. #AP6514d) IHC analysis in formalin fixed and paraffin embedded human tonsil tissue followed by peroxidase conjugation of the secondary antibody and DAB

staining. This data demonstrates the use of the FLI-1 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

#### **FLI1 Antibody (N-term K67) - Background**

FLI1 is a sequence-specific transcriptional activator. It recognizes the DNA sequence 5'-C[CA]GGAAGT-3'.

#### **FLI1 Antibody (N-term K67) - References**

Embree,L.J., Cancer Res. 69 (10), 4363-4371 (2009)  
Torlakovic,E.E., Histol. Histopathol. 23 (11), 1309-1314 (2008)  
Cironi,L., PLoS ONE 3 (7), E2634 (2008)