

**WNT4 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP6683B****Specification**

---

**WNT4 Antibody (C-term) - Product Information**

|                   |   |
|-------------------|---|
| Application       | WB, IHC-P,E                                     |
| Primary Accession | <a href="#">P56705</a>                          |
| Other Accession   | <a href="#">Q9OXQ5</a> , <a href="#">P22724</a> |
| Reactivity        | Human   |
| Predicted         | Mouse, Rat                                      |
| Host              | Rabbit  |
| Clonality         | Polyclonal                                      |
| Isotype           | Rabbit IgG                                      |
| Antigen Region    | 242-269   |

**WNT4 Antibody (C-term) - Additional Information****Gene ID** 54361**Other Names**

Protein Wnt-4, WNT4

**Target/Specificity**

This WNT4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 242-269 amino acids from the C-terminal region of human WNT4.

**Dilution**

WB~~1:1000

IHC-P~~1:50~100

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

WNT4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**WNT4 Antibody (C-term) - Protein Information****Name** WNT4**Function** Ligand for members of the frizzled family of seven transmembrane receptors (Probable).

Plays an important role in the embryonic development of the urogenital tract and the lung (PubMed:[15317892](#), PubMed:[16959810](#), PubMed:[18179883](#), PubMed:[18182450](#)). Required for normal mesenchyme to epithelium transition during embryonic kidney development. Required for the formation of early epithelial renal vesicles during kidney development (By similarity). Required for normal formation of the Mullerian duct in females, and normal levels of oocytes in the ovaries (PubMed:[15317892](#), PubMed:[16959810](#), PubMed:[18182450](#)). Required for normal down-regulation of 3 beta-hydroxysteroid dehydrogenase in the ovary (PubMed:[15317892](#), PubMed:[16959810](#), PubMed:[18182450](#)). Required for normal lung development and for normal patterning of tracheal cartilage rings (By similarity).

#### **Cellular Location**


Secreted, extracellular space, extracellular matrix

#### **WNT4 Antibody (C-term) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **WNT4 Antibody (C-term) - Images**



Western blot analysis of WNT4 antibody (C-term) (Cat. #AP6683b) in 293 cell line lysates (35ug/lane). WNT4 (arrow) was detected using the purified Pab.



(LEFT) Formalin-fixed and paraffin-embedded human lung carcinoma reacted with WNT4 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. (RIGHT) Flow cytometric analysis of 293 cells using WNT4 Antibody (C-term) (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### WNT4 Antibody (C-term) - Background

The WNT family consists of secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. WNT4 is a protein which shows 98% amino acid identity to the Wnt4 protein of mouse and rat. It plays a concerted role in both the control of female development and the prevention of testes formation.

### WNT4 Antibody (C-term) - References

Kuulasmaa, T., Horm. Metab. Res. 40 (10), 668-673 (2008) Miyakoshi, T., Endocr. Pathol. 19 (4), 261-273 (2008)

### WNT4 Antibody (C-term) - Citations

- [β-catenin is a central mediator of pro-fibrotic Wnt signaling in systemic sclerosis.](#)