

CYP2C19 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6710c**Specification**

CYP2C19 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	P33261
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	257-285

CYP2C19 Antibody (Center) - Additional Information**Gene ID** 1557**Other Names**

Cytochrome P450 2C19, 11413-, (R)-limonene 6-monooxygenase, (S)-limonene 6-monooxygenase, (S)-limonene 7-monooxygenase, CYP11C17, CYP11C19, Cytochrome P450-11A, Cytochrome P450-254C, Mephenytoin 4-hydroxylase, CYP2C19

Target/Specificity

This CYP2C19 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 257-285 amino acids from the Central region of human CYP2C19.

Dilution

WB~~1:1000

IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CYP2C19 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

CYP2C19 Antibody (Center) - Protein Information**Name** CYP2C19**Function** A cytochrome P450 monooxygenase involved in the metabolism of polyunsaturated

fatty acids (PUFA) (PubMed:[18577768](#), PubMed:[19965576](#), PubMed:[20972997](#)). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:[18577768](#), PubMed:[19965576](#), PubMed:[20972997](#)). Catalyzes the hydroxylation of carbon-hydrogen bonds. Hydroxylates PUFA specifically at the omega-1 position (PubMed:[18577768](#)). Catalyzes the epoxidation of double bonds of PUFA (PubMed:[20972997](#), PubMed:[19965576](#)). Also metabolizes plant monoterpenes such as limonene. Oxygenates (R)- and (S)-limonene to produce carveol and perillyl alcohol (PubMed:[11950794](#)). Responsible for the metabolism of a number of therapeutic agents such as the anticonvulsant drug S-mephenytoin, omeprazole, proguanil, certain barbiturates, diazepam, propranolol, citalopram and imipramine. Hydroxylates fenbendazole at the 4' position (PubMed:[23959307](#)).

Cellular Location


Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein

CYP2C19 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CYP2C19 Antibody (Center) - Images



Anti-CYP2C19 Antibody (Center) at 1:2000 dilution + human liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 56 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



Western blot analysis of CYP2C19 Antibody (Center) (Cat. #AP6710c) in Jurkat cell line lysates (35ug/lane). CYP2C19 (arrow) was detected using the purified Pab.



CYP2C19 Antibody (Center) (Cat. #AP6710c) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of CYP2C19 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

CYP2C19 Antibody (Center) - Background

CYP2C19 is a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and is known to metabolize many xenobiotics, including the anticonvulsive drug mephenytoin, omeprazole, diazepam and some barbiturates. Polymorphism within its gene is associated with variable ability to metabolize mephenytoin, known as the poor metabolizer and extensive metabolizer phenotypes.

CYP2C19 Antibody (Center) - References

Shuldiner, A.R., JAMA 302 (8), 849-857 (2009)
Nelson, D.R., Pharmacogenetics 14 (1), 1-18 (2004)