

HPRT1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP6732B

Specification

HPRT1 Antibody (C-term) - Product Information

Primary AccessionOther AccessionReactivityPredictedHostClonality	WB, IHC-P, FC,E <u>P00492</u> <u>Q6LDD9</u> Human Monkey Rabbit Polyclonal Rabbit IgG
	_
Antigen Region	150-178

HPRT1 Antibody (C-term) - Additional Information

Gene ID 3251

Other Names Hypoxanthine-guanine phosphoribosyltransferase, HGPRT, HGPRTase, HPRT1, HPRT

Target/Specificity This HPRT1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 150-178 amino acids from the C-terminal region of human HPRT1.

Dilution WB~~1:1000 IHC-P~~1:10~50 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HPRT1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

HPRT1 Antibody (C-term) - Protein Information

Name HPRT1



Synonyms HPRT

Function Converts guanine to guanosine monophosphate, and hypoxanthine to inosine monophosphate. Transfers the 5-phosphoribosyl group from 5- phosphoribosylpyrophosphate onto the purine. Plays a central role in the generation of purine nucleotides through the purine salvage pathway.


Cellular Location Cytoplasm.

HPRT1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

HPRT1 Antibody (C-term) - Images




Western blot analysis of HPRT1 antibody (C-term) (Cat. #AP6732b) in Hela cell line lysates (35ug/lane). HPRT1 (arrow) was detected using the purified Pab.






Western blot analysis of HPRT1 (arrow) using rabbit polyclonal HPRT1 Antibody (C-term) (Cat. #AP6732b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the HPRT1 gene.



HPRT1 Antibody (C-term) (Cat. #AP6732b) IHC analysis in formalin fixed and paraffin embedded mouse brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the HPRT1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.





HPRT1 Antibody (C-term) (Cat. #AP6732b) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

HPRT1 Antibody (C-term) - Background

The protein is a transferase, which catalyzes conversion of hypoxanthine to inosine monophosphate and guanine to guanosine monophosphate via transfer of the 5-phosphoribosyl group from 5-phosphoribosyl 1-pyrophosphate. This enzyme plays a central role in the generation of purine nucleotides through the purine salvage pathway.

HPRT1 Antibody (C-term) - References

Ceballos-Picot, I., Hum. Mol. Genet. 18 (13), 2317-2327 (2009) Sculley, D.G., Hum. Genet. 90 (3), 195-207 (1992)