

GBAS Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP6752c

Specification

GBAS Antibody (Center) - Product Information

Application WB, IHC-P, FC,E

Primary Accession
Reactivity
Host
Clonality
Isotype
Calculated MW
Antigen Region

O75323
Human
Rabbit
Polyclonal
Rabbit IgG
33743
Antigen Region
130-159

GBAS Antibody (Center) - Additional Information

Gene ID 2631

Other Names

Protein NipSnap homolog 2, NipSnap2, Glioblastoma-amplified sequence, GBAS, NIPSNAP2

Target/Specificity

This GBAS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 130-159 amino acids from the Central region of human GBAS.

Dilution

WB~~1:1000 IHC-P~~1:50~100 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GBAS Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

GBAS Antibody (Center) - Protein Information

Name NIPSNAP2 (HGNC:4179)

Synonyms GBAS



Function May act as a positive regulator of L-type calcium channels.

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:055126}. Mitochondrion outer membrane

Tissue Location


Widely expressed. Most abundant in heart and skeletal muscle

GBAS Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

GBAS Antibody (Center) - Images



Western blot analysis of GBAS Antibody (Center) (Cat. #AP6752c) in Ramos cell line lysates (35ug/lane).GBAS (arrow) was detected using the purified Pab.




GBAS Antibody (Center) (Cat. #AP6752c) IHC analysis in formalin fixed and paraffin embedded



relevance has not been evaluated.

skeletal muscle followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the GBAS Antibody (Center) for immunohistochemistry. Clinical



GBAS Antibody (Center) (Cat. #AP6752c) flow cytometric analysis of Ramos cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

GBAS Antibody (Center) - Background

Chromosomal region 7p12, which contains GBAS, is amplified in approximately 40% of glioblastomas, the most common and malignant form of central nervous system tumor. The predicted 286-amino acid protein contains a signal peptide, a transmembrane domain, and 2 tyrosine phosphorylation sites. The GBAS transcript is expressed most abundantly in heart and skeletal muscle. GBAS protein might be involved in vesicular transport.

GBAS Antibody (Center) - References

Simpson, J.C., et al. EMBO Rep. 1(3):287-292(2000) Seroussi, E., et al. Gene 212(1):13-20(1998) Wang, X.Y., et al. Genomics 49(3):448-451(1998) GBAS Antibody (Center) - Citations

• NIPSNAP1 and NIPSNAP2 Act as