

**FMO3 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP6901a****Specification**

---

**FMO3 Antibody (N-term) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">P31513</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	60033
Antigen Region	30-56

**FMO3 Antibody (N-term) - Additional Information****Gene ID** 2328**Other Names**

Dimethylaniline monooxygenase [N-oxide-forming] 3, Dimethylaniline oxidase 3, FMO II, FMO form 2, Hepatic flavin-containing monooxygenase 3, FMO 3, Trimethylamine monooxygenase, FMO3

**Target/Specificity**

This FMO3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 30-56 amino acids from the N-terminal region of human FMO3.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

FMO3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**FMO3 Antibody (N-term) - Protein Information****Name** FMO3

**Function** Essential hepatic enzyme that catalyzes the oxygenation of a wide variety of nitrogen- and sulfur-containing compounds including drugs as well as dietary compounds (PubMed:[10759686](#), PubMed:[30381441](#), PubMed:[32156684](#)). Plays an important role in the metabolism of trimethylamine (TMA), via the production of trimethylamine N-oxide (TMAO) metabolite (PubMed:[9776311](#)). TMA is generated by the action of gut microbiota using dietary precursors such as choline, choline containing compounds, betaine or L-carnitine. By regulating TMAO concentration, FMO3 directly impacts both platelet responsiveness and rate of thrombus formation (PubMed:[29981269](#)).

#### Cellular Location

Microsome membrane {ECO:0000250|UniProtKB:P32417}; Single-pass membrane protein.  
Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:P32417}; Single-pass membrane protein

#### Tissue Location


Liver.

### FMO3 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### FMO3 Antibody (N-term) - Images



Western blot analysis of FMO3 Antibody (N-term) (Cat. #AP6901a) in MDA-MB231 cell line lysates (35ug/lane). FMO3 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human hepatocarcinoma reacted with FMO3 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



FMO3 Antibody (N-term) (Cat. #AP6901a) flow cytometric analysis of MDA-MB231 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### FMO3 Antibody (N-term) - Background

FMO3 is involved in the oxidative metabolism of a variety of xenobiotics such as drugs and pesticides. It N-oxygenates primary aliphatic alkylamines as well as secondary and tertiary amines. It plays an important role in the metabolism of trimethylamine (TMA), via the production of TMA N-oxide (TMAO). Is also able to perform S-oxidation when acting on sulfide compounds.

### FMO3 Antibody (N-term) - References

Allerston, C.K., et.al., Mol. Genet. Metab. 98 (1-2), 198-202 (2009)