10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# GRK4 Antibody (C-term)

**Purified Rabbit Polyclonal Antibody (Pab)**
**Catalog # AP7006a**

## Specification

### GRK4 Antibody (C-term) - Product Information

| | |
|---|---|
| Application | **WB, IHC-P,E** |
| Primary Accession | P32298 |
| Reactivity | **Human** |
| Host | **Rabbit** |
| Clonality | **Polyclonal** |
| Isotype | **Rabbit IgG** |
| Antigen Region | **549-577** |

### GRK4 Antibody (C-term) - Additional Information

**Gene ID** 2868

**Other Names**
G protein-coupled receptor kinase 4, G protein-coupled receptor kinase GRK4, ITI1, GRK4, GPRK2L, GPRK4

**Target/Specificity**
This GRK4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 549-577 amino acids from the C-terminal region of human GRK4.

**Dilution**
WB~~1:1000
IHC-P~~1:10~50

**Format**
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**
GRK4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### GRK4 Antibody (C-term) - Protein Information

**Name** GRK4

**Synonyms** GPRK2L, GPRK4

**Function** Specifically phosphorylates the activated forms of G protein- coupled receptors. GRK4-alpha can phosphorylate rhodopsin and its activity is inhibited by calmodulin; the other three isoforms do not phosphorylate rhodopsin and do not interact with calmodulin. GRK4-alpha and GRK4-gamma phosphorylate DRD3. Phosphorylates ADRB2.

**Cellular Location**
Cytoplasm. Cytoplasm, cell cortex. Note=Both at the cell surface and dispersed in the cytoplasm under basal conditions Receptor stimulation results in the internalization of GRK4 to the perinuclear area, where colocalization with DRD3 is observed strongly at 5 and 15 minutes. DRD3 and GRK4 colocalize in lipid rafts of renal proximal tubule cells


**Tissue Location**
Isoform 1, isoform 2, isoform 3, and isoform 4 are expressed in testis. Isoform 4 is expressed in myometrium

## GRK4 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## GRK4 Antibody (C-term) - Images




Western blot analysis of anti-GRK4 Pab (Cat. #AP7006a) in HL60 cell lysate. GRK4 (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999




Formalin-fixed and paraffin-embedded human testis tissue reacted with GRK4 antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

**GRK4 Antibody (C-term) - Background**

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the g phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The AGC kinase group consists of 63 kinases including the cyclic nucleotide-regulated protein kinase (PKA & PKG) family, the diacylglycerol-activated/phospholipid-dependent protein kinase C (PKC) family, the related to PKA and PKC (RAC/Akt) protein kinase family, the kinases that phosphorylate G protein-coupled receptors family (ARK), and the kinases that phosphorylate ribosomal protein S6 family (RSK).

**GRK4 Antibody (C-term) - References**

Perroy, J., et al., EMBO J. 22(15):3816-3824 (2003).
Watanabe, H., et al., Kidney Int. 62(3):790-798 (2002).
Sallese, M., et al., J. Biol. Chem. 272(15):10188-10195 (1997).
Premont, R.T., et al., J. Biol. Chem. 271(11):6403-6410 (1996).
Sallese, M., et al., Biochem. Biophys. Res. Commun. 199(2):848-854 (1994).

**GRK4 Antibody (C-term) - Citations**

- G protein-coupled receptor kinase 4 (GRK4) regulates the phosphorylation and function of the dopamine D3 receptor.