

**CKB Antibody (N-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP7059a****Specification**

---

**CKB Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P12277</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	1-30

**CKB Antibody (N-term) - Additional Information****Gene ID** 1152**Other Names**

Creatine kinase B-type, B-CK, Creatine kinase B chain, CKB, CKBB

**Target/Specificity**

This CKB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human CKB.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CKB Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CKB Antibody (N-term) - Protein Information****Name** CKB ([HGNC:1991](#))**Synonyms** CKBB

**Function** Reversibly catalyzes the transfer of phosphate between ATP and various phosphogens (e.g. creatine phosphate) (PubMed:[8186255](#)). Creatine kinase isoenzymes play a central role in

energy transduction in tissues with large, fluctuating energy demands, such as skeletal muscle, heart, brain and spermatozoa (Probable). Acts as a key regulator of adaptive thermogenesis as part of the futile creatine cycle: localizes to the mitochondria of thermogenic fat cells and acts by mediating phosphorylation of creatine to initiate a futile cycle of creatine phosphorylation and dephosphorylation (By similarity). During the futile creatine cycle, creatine and N-phosphocreatine are in a futile cycle, which dissipates the high energy charge of N- phosphocreatine as heat without performing any mechanical or chemical work (By similarity).

#### Cellular Location


Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q04447}. Mitochondrion {ECO:0000250|UniProtKB:Q04447}. Cell membrane. Note=Localizes to the mitochondria of thermogenic fat cells via the internal MTS-like signal (iMTS-L) region {ECO:0000250|UniProtKB:Q04447}

#### CKB Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### CKB Antibody (N-term) - Images



The anti-CKB Pab (Cat. #AP7059a) is used in Western blot to detect CKB in mouse colon tissue lysate (Lane 1) and Y79 cell lysate (Lane 2).



Western blot analysis of CKB (arrow) using rabbit polyclonal CKB Antibody (N5) (Cat. #AP7059a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the CKB gene.

#### **CKB Antibody (N-term) - Background**

Creatine kinase isoenzymes play a central role in energy transduction in tissues with large, fluctuating energy demands, such as skeletal muscle, heart, brain and spermatozoa. The CKB isoform, present in many tissues but especially brain, is a cytoplasmic enzyme involved in energy homeostasis. CKB reversibly catalyzes the transfer of phosphate between ATP and various phosphogens such as creatine phosphate. Creatine kinase B-driven energy transfer in the brain is important for habituation and spatial learning behaviour, mossy fibre field size and determination of seizure susceptibility. The encoded protein is a member of the ATP:guanido phosphotransferase protein family.