

Ethanolamine Kinase (EKI1) Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7068b

Specification

Ethanolamine Kinase (EKI1) Antibody (C-term) - Product Information

Application WB,E
Primary Accession Q9HBU6

Reactivity Human, Hamster, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 50968
Antigen Region 356-386

Ethanolamine Kinase (EKI1) Antibody (C-term) - Additional Information

Gene ID 55500

Other Names

Ethanolamine kinase 1, EKI 1, ETNK1, EKI1

Target/Specificity

This Ethanolamine Kinase (EKI1) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 356-386 amino acids from the C-terminal region of human Ethanolamine Kinase (EKI1).

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Ethanolamine Kinase (EKI1) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Ethanolamine Kinase (EKI1) Antibody (C-term) - Protein Information

Name ETNK1 (HGNC:24649)

Function Highly specific for ethanolamine phosphorylation. May be a rate-controlling step in phosphatidylethanolamine biosynthesis.



Cellular Location Cytoplasm.

Tissue Location


Expressed in kidney, liver, placenta, heart, leukocyte, ovary and testis.

Ethanolamine Kinase (EKI1) Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Ethanolamine Kinase (EKI1) Antibody (C-term) - Images



The anti-EKI1 Pab (Cat. #AP7068b) is used in Western blot to detect EKI1 in CHO cell lysate.

Ethanolamine Kinase (EKI1) Antibody (C-term) - Background

Ethanolamine kinase 1 (EKI1), functions in the first committed step of the phosphatidylethanolamine synthesis pathway. This cytosolic enzyme is specific for ethanolamine and exhibits negligible kinase activity on choline.