

STK38 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7074b**Specification**

STK38 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q15208
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	410-439

STK38 Antibody (C-term) - Additional Information**Gene ID** 11329**Other Names**

Serine/threonine-protein kinase 38, NDR1 protein kinase, Nuclear Dbf2-related kinase 1, STK38
{ECO:0000312|EMBL:AAH120851}

Target/Specificity

This STK38 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 410-439 amino acids from the C-terminal region of human STK38.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

STK38 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

STK38 Antibody (C-term) - Protein Information**Name** STK38 {ECO:0000312|EMBL:AAH12085.1}

Function Negative regulator of MAP3K1/2 signaling. Converts MAP3K2 from its phosphorylated form to its non-phosphorylated form and inhibits autophosphorylation of MAP3K2.

Cellular Location

Nucleus. Cytoplasm.


Tissue Location

Ubiquitously expressed with highest levels observed in peripheral blood leukocytes.

STK38 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STK38 Antibody (C-term) - Images

The anti-STK38 Pab (Cat. #AP7074b) is used in Western blot to detect STK38 in SK-Br-3 (left) and Jurkat (right) cell line lysates.

STK38 Antibody (C-term) - Background

Stk38, also known as NUCLEAR DBF2-RELATED PROTEIN, contains 1 protein kinase domain that interacts with mob1 and mob2. The homodimeric s100b binds to two molecules of stk38. It is ubiquitously expressed with highest levels observed in peripheral blood leukocytes. Stk38 is activated by binding of s100b which releases autoinhibitory n-lobe interactions, enabling atp to bind and the autophosphorylation of ser-281. thr-444 then undergoes calcium-dependent phosphorylation by an upstream kinase. Interactions between phosphorylated thr-444 and the n-lobe promote additional structural changes that complete the activation of the kinase. Autoinhibition is also released by the binding of mob1/mobk11a and mob2/hcca2 to the n-terminal of stk38.

STK38 Antibody (C-term) - References

Devroe, E., et al., J. Biol. Chem. 279(23):24444-24451 (2004).

Tamaskovic, R., et al., J. Biol. Chem. 278(9):6710-6718 (2003).
Tripodis, N., et al., Genome Res. 8(6):631-643 (1998).
Millward, T., et al., Proc. Natl. Acad. Sci. U.S.A. 92(11):5022-5026 (1995).