

**CLASP2 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP7181b**

**Specification**

---

**CLASP2 Antibody (C-term) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">O75122</a>
Other Accession	<a href="#">NP_055912</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	141064
Antigen Region	946-975

**CLASP2 Antibody (C-term) - Additional Information**

**Gene ID** 23122

**Other Names**

CLIP-associating protein 2, Cytoplasmic linker-associated protein 2, Protein Orbit homolog 2, hOrbit2, CLASP2, KIAA0627

**Target/Specificity**

This CLASP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 946-975 amino acids from the C-terminal region of human CLASP2.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CLASP2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CLASP2 Antibody (C-term) - Protein Information**

**Name** CLASP2

**Synonyms** KIAA0627

**Function** Microtubule plus-end tracking protein that promotes the stabilization of dynamic microtubules (PubMed:[26003921](#)). Involved in the nucleation of noncentrosomal microtubules originating from the trans-Golgi network (TGN). Required for the polarization of the cytoplasmic microtubule arrays in migrating cells towards the leading edge of the cell. May act at the cell cortex to enhance the frequency of rescue of depolymerizing microtubules by attaching their plus-ends to cortical platforms composed of ERC1 and PHLDB2 (PubMed:[16824950](#)). This cortical microtubule stabilizing activity is regulated at least in part by phosphatidylinositol 3-kinase signaling. Also performs a similar stabilizing function at the kinetochore which is essential for the bipolar alignment of chromosomes on the mitotic spindle (PubMed:[16866869](#), PubMed:[16914514](#)). Acts as a mediator of ERBB2- dependent stabilization of microtubules at the cell cortex.

**Cellular Location**

Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Chromosome, centromere, kinetochore. Cytoplasm, cytoskeleton, spindle. Golgi apparatus {ECO:0000250|UniProtKB:Q8BRT1}. Golgi apparatus, trans-Golgi network. Cell membrane. Cell projection, ruffle membrane. Note=Localizes to microtubule plus ends (PubMed:15631994). Localizes to centrosomes, kinetochores and the mitotic spindle from prometaphase. Subsequently localizes to the spindle midzone from anaphase and to the midbody from telophase (PubMed:16866869, PubMed:16914514). In migrating cells localizes to the plus ends of microtubules within the cell body and to the entire microtubule lattice within the lamella. Localizes to the cell cortex and this requires ERC1 and PHLDB2 (PubMed:16824950). The MEMO1-RHOA-DIAPH1 signaling pathway controls localization of the phosphorylated form to the cell membrane.

**Tissue Location**


Brain-specific.

**CLASP2 Antibody (C-term) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CLASP2 Antibody (C-term) - Images**



Western blot analysis of anti-CLASP2 Antibody (C-term) (Cat. #AP7181b) in mouse kidney tissue lysates (35ug/lane). CLASP2(arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human Hepatocellular carcinoma reacted with anti-CLASP2 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

#### **CLASP2 Antibody (C-term) - Background**

Microtubule plus-end tracking protein that promotes the stabilization of dynamic microtubules. Required for the polarization of the cytoplasmic microtubule arrays in migrating cells towards the leading edge of the cell. May act at the cell cortex to enhance the frequency of rescue of depolymerizing microtubules by attaching their plus-ends to cortical platforms composed of ERC1 and PHLDB2. This cortical microtubule stabilizing activity is regulated at least in part by phosphatidylinositol 3-kinase signaling. Also performs a similar stabilizing function at the kinetochore which is essential for the bipolar alignment of chromosomes on the mitotic spindle.

#### **CLASP2 Antibody (C-term) - References**

Pereira,A.L., Mol. Biol. Cell 17 (10), 4526-4542 (2006)  
Mimori-Kiyosue,Y., Genes Cells 11 (8), 845-857 (2006)  
Lansbergen,G., Dev. Cell 11 (1), 21-32 (2006)