

DRAK2 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7221b**Specification**

DRAK2 Antibody (C-term) - Product Information

Application	IF, WB, IHC-P,E
Primary Accession	O94768
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	342-371

DRAK2 Antibody (C-term) - Additional Information**Gene ID** 9262**Other Names**

Serine/threonine-protein kinase 17B, DAP kinase-related apoptosis-inducing protein kinase 2, STK17B, DRAK2

Target/Specificity

This DRAK2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 342-371 amino acids from the C-terminal region of human DRAK2.

Dilution

IF~~1:10~50
WB~~1:1000
IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DRAK2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DRAK2 Antibody (C-term) - Protein Information**Name** STK17B**Synonyms** DRAK2

Function Phosphorylates myosin light chains (By similarity). Acts as a positive regulator of apoptosis.

Cellular Location

Nucleus. Cell membrane. Endoplasmic reticulum-Golgi intermediate compartment.
Note=Colocalizes with STK17B at the plasma membrane.

Tissue Location


Highly expressed in placenta, lung, pancreas. Lower levels in heart, brain, liver, skeletal muscle and kidney

DRAK2 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


DRAK2 Antibody (C-term) - Images



IF staining of Drak2 in WT versus Tg T cells. Permeabilized lymph node T cells were double-stained with rabbit anti-Drak2/sheep anti-rabbit Ig-FITC and anti-Thy1.2-PE monoclonal antibody, and signals were registered by confocal microscopy. THE cell surface Thy1.2 is shown in red, and intracellular Drak2 is in green.



The anti-DRAK2 Pab (Cat. #AP7221b) is used in Western blot to detect DRAK2 in mouse lung tissue lysate.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

DRAK2 Antibody (C-term) - Background

DRAK2 is a novel serine/threonine kinase that induces apoptosis via catalytic activity. DRAKs present high sequence homology to DAP and ZIP kinases, and they represent a novel family of serine/threonine kinases. DRAK2 is located in nucleus, and the messenger RNA is ubiquitously expressed in human tissues.

DRAK2 Antibody (C-term) - References

Sanjo, H., et al., J. Biol. Chem. 273(44):29066-29071 (1998).

DRAK2 Antibody (C-term) - Citations

- [Drak2 is upstream of p70S6 kinase: its implication in cytokine-induced islet apoptosis, diabetes, and islet transplantation.](#)
- [Transgenic drak2 overexpression in mice leads to increased T cell apoptosis and compromised memory T cell development.](#)