

PIST Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7262b

Specification

PIST Antibody (C-term) - Product Information

Application WB, IHC-P,E
Primary Accession O9HD26
Other Accession O8BH60

Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Antigen Region 361-390

PIST Antibody (C-term) - Additional Information

Gene ID 57120

Other Names

Golgi-associated PDZ and coiled-coil motif-containing protein, CFTR-associated ligand, Fused in glioblastoma, PDZ protein interacting specifically with TC10, PIST, GOPC, CAL, FIG

Target/Specificity

This PIST antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 361-390 amino acids from the C-terminal region of human PIST.

Dilution

WB~~1:1000 IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PIST Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PIST Antibody (C-term) - Protein Information

Name GOPC (HGNC:17643)

Function Plays a role in intracellular protein trafficking and degradation (PubMed: 11707463,



Tel: 858.875.1900 Fax: 858.875.1999

PubMed:14570915, PubMed:15358775). May regulate CFTR chloride currents and acid-induced ASIC3 currents by modulating cell surface expression of both channels (By similarity). May also regulate the intracellular trafficking of the ADR1B receptor (PubMed:15358775). May play a role in autophagy (By similarity). Together with MARCHF2 mediates the ubiquitination and lysosomal degradation of CFTR (PubMed: 23818989). Overexpression results in CFTR intracellular retention and lysosomaldegradation in the lysosomes (PubMed: 11707463, PubMed: 14570915).

Cellular Location

Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein. Golgi apparatus, trans-Golgi network membrane; Peripheral membrane protein Synapse. Postsynaptic density. Cell projection, dendrite. Note=Enriched in synaptosomal and postsynaptic densities (PSD) fractions. Expressed in cell bodies and dendrites of Purkinje cells. Localized at the trans-Golgi network (TGN) of spermatids and the medulla of round spermatides.

Tissue Location


Ubiquitously expressed.

PIST Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

PIST Antibody (C-term) - Images



Western blot analysis of PIST Antibody (C-term) (Cat.#AP7262b) in Ramos cell line lysates (35ug/lane). PIST (arrow) was detected using the purified Pab.





Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with PIST antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

PIST Antibody (C-term) - Background

PDZ domains contain approximately 90 amino acids and bind the extreme C terminus of proteins in a sequence-specific manner. PIST, a PDZ domain-containing Golgi protein, was discovered in a yeast two-hybrid system as a binding partner to Beclin-1, a Bcl-2-interacting protein homologous to the yeast autophagy gene apg6. Experiments with mutant PIST proteins lacking the PDZ domain showed that PIST interaction with Beclin-1 through its coiled-coil domain can modulate Beclin-1 activity and suggest that PIST interactions with other proteins through its PDZ domain may regulate the activity of PIST and Beclin-1.

PIST Antibody (C-term) - References

Li,X., Protein Sci. 15 (9), 2149-2158 (2006) Ito,H., Biochem. J. 397 (3), 389-398 (2006) Wente,W., J. Biol. Chem. 280 (37), 32419-32425 (2005)

PIST Antibody (C-term) - Citations

• A rapid method to improve protein detection by indirect ELISA.