

MFAP5 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7319b

Specification

MFAP5 Antibody (C-term) - Product Information

Application WB, IHC-P, FC,E

Primary Accession <u>Q13361</u>

Other Accession <u>Q9QZJ6</u>, <u>Q28022</u>

Reactivity Human

Predicted Bovine, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 19612
Antigen Region 133-160

MFAP5 Antibody (C-term) - Additional Information

Gene ID 8076

Other Names

Microfibrillar-associated protein 5, MFAP-5, MP25, Microfibril-associated glycoprotein 2, MAGP-2, MFAP5, MAGP2

Target/Specificity

This MFAP5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 133-160 amino acids from the C-terminal region of human MFAP5.

Dilution

WB~~1:1000 IHC-P~~1:50~100 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.


Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MFAP5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MFAP5 Antibody (C-term) - Protein Information





Name MFAP5

Synonyms MAGP2

Function May play a role in hematopoiesis. In the cardiovascular system, could regulate growth factors or participate in cell signaling in maintaining large vessel integrity (By similarity). Component of the elastin-associated microfibrils (PubMed:8557636).

Cellular Location


Secreted, extracellular space, extracellular matrix

MFAP5 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

MFAP5 Antibody (C-term) - Images




Western blot analysis of MFAP5 antibody (C-term) (Cat.#AP7319b) in K562 cell line lysates (35ug/lane). MFAP5 (arrow) was detected using the purified Pab.






Western blot analysis of MFAP5 (arrow) using rabbit polyclonal MFAP5 Antibody (C-term) (Cat. #AP7319b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the MFAP5 gene.




Formalin-fixed and paraffin-embedded human Skeletal muscle reacted with MFAP5 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



MFAP5 Antibody (C-term) (Cat. #AP7319b) flow cytometry analysis of K562 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

MFAP5 Antibody (C-term) - Background

MFAP5 is a 25-kD microfibril-associated glycoprotein which is rich in serine and threonine residues.





Tel: 858.875.1900 Fax: 858.875.1999

The protein lacks a hydrophobic carboxyl terminus and proline-, glutamine-, and tyrosine-rich regions, which are characteristics of a related 31-kDa microfibril-associated glycoprotein (MFAP2). The close similarity between these two proteins is confined to a central region of 60 aa where precise alignment of 7 cysteine residues occurs. The structural differences suggest that this protein has some functions that are distinct from those of MFAP2.

MFAP5 Antibody (C-term) - References

Albig, A.R., Becenti, D.J. Microvasc. Res. 76 (1), 7-14 (2008) Miyamoto, A., Lau, R. J. Biol. Chem. 281 (15), 10089-10097 (2006) Penner, A.S., Rock, M.J. J. Biol. Chem. 277 (38), 35044-35049 (2002) Hatzinikolas, G. J. Biol. Chem. 273 (45), 29309-29314 (1998)