

S100A6 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7346b**Specification**

S100A6 Antibody (C-term) - Product Information

Application	WB, FC,E
Primary Accession	P06703
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	10180
Antigen Region	37-67

S100A6 Antibody (C-term) - Additional Information**Gene ID** 6277**Other Names**

Protein S100-A6, Calcyclin, Growth factor-inducible protein 2A9, MLN 4, Prolactin receptor-associated protein, PRA, S100 calcium-binding protein A6, S100A6, CACY

Target/Specificity

This S100A6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 37-67 amino acids from the C-terminal region of human S100A6.

Dilution

WB~~1:1000

FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

S100A6 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

S100A6 Antibody (C-term) - Protein Information**Name** S100A6**Synonyms** CACY

Function May function as calcium sensor and modulator, contributing to cellular calcium signaling. May function by interacting with other proteins, such as TPR-containing proteins, and indirectly play a role in many physiological processes such as the reorganization of the actin cytoskeleton and in cell motility. Binds 2 calcium ions. Calcium binding is cooperative.

Cellular Location


Nucleus envelope. Cytoplasm. Cell membrane; Peripheral membrane protein; Cytoplasmic side

S100A6 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


S100A6 Antibody (C-term) - Images



Western blot analysis of S100A6 antibody (C-term) (Cat.#AP7346b) in mouse lung tissue lysates (35ug/lane). S100A6 (arrow) was detected using the purified Pab.



Western blot analysis of S100A6 (arrow) using rabbit polyclonal S100A6 Antibody (C-term) (Cat. #AP7346b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the S100A6 gene (Lane 2) (Origene Technologies).



S100A6 Antibody (C-term) (Cat. #AP7346b) flow cytometric analysis of HeLa cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

S100A6 Antibody (C-term) - Background

S100A6 is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. This protein may function in stimulation of Ca²⁺-dependent insulin release, stimulation of prolactin secretion, and exocytosis.

S100A6 Antibody (C-term) - References

- van Dieck, J. J. Biol. Chem. 284 (20), 13804-13811 (2009)
- Tsoporis, J.N., Izhar, S. J. Biol. Chem. 283 (44), 30174-30183 (2008)
- Lee, Y.T., Dimitrova, Y.N. Biochemistry 47 (41), 10921-10932 (2008)
- Melle, C., Ernst, G. PLoS ONE 3 (12), E3767 (2008)