

FBP1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7385c

Specification

FBP1 Antibody (Center) - Product Information

Application WB, IHC-P,E
Primary Accession P09467
Other Accession Q90XD6

Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 36842
Antigen Region 125-156

FBP1 Antibody (Center) - Additional Information

Gene ID 2203

Other Names

Fructose-1, 6-bisphosphatase 1, FBPase 1, D-fructose-1, 6-bisphosphate 1-phosphohydrolase 1, Liver FBPase, FBP1, FBP

Target/Specificity

This FBP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 125-156 amino acids from the Central region of human FBP1.

Dilution

WB~~1:1000 IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FBP1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

FBP1 Antibody (Center) - Protein Information

Name FBP1





Synonyms FBP

Function Catalyzes the hydrolysis of fructose 1,6-bisphosphate to fructose 6-phosphate in the presence of divalent cations, acting as a rate-limiting enzyme in gluconeogenesis. Plays a role in regulating glucose sensing and insulin secretion of pancreatic beta-cells. Appears to modulate glycerol gluconeogenesis in liver. Important regulator of appetite and adiposity; increased expression of the protein in liver after nutrient excess increases circulating satiety hormones and reduces appetite-stimulating neuropeptides and thus seems to provide a feedback mechanism to limit weight gain.

Tissue Location


Expressed in pancreatic islets.

FBP1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

FBP1 Antibody (Center) - Images




Western blot analysis of FBP1 Antibody (Center) (Cat.# AP7385c) in MCF-7 cell line lysates (35ug/lane). FBP1 (arrow) was detected using the purified Pab.






Western blot analysis of FBP1 antibody (Center) (Cat.# AP7385c) in mouse kidney tissue lysates (35ug/lane). FBP1 (arrow) was detected using the purified Pab.



Western blot analysis of FBP1 (arrow) using rabbit polyclonal FBP1 Antibody (Center) (Cat. #AP7385c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the FBP1 gene.




FBP1 Antibody (Center) (Cat.# AP7385c) IHC analysis in formalin fixed and paraffin embedded human hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the FBP1 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

FBP1 Antibody (Center) - Background

Fructose-1,6-bisphosphatase 1, a gluconeogenesis regulatory enzyme, catalyzes the hydrolysis of





fructose 1,6-bisphosphate to fructose 6-phosphate and inorganic phosphate. Fructose-1,6-diphosphatase deficiency is associated with hypoglycemia and metabolic acidosis.

FBP1 Antibody (Center) - References

Visinoni, S., Am. J. Physiol. Endocrinol. Metab. 295 (5), E1132-E1141 (2008) Kebede, M., Diabetes 57 (7), 1887-1895 (2008)