

DDX5 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7459b**Specification**

DDX5 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P17844
Other Accession	Q61656 , Q4R6M5
Reactivity	Human
Predicted	Monkey, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	69148
Antigen Region	491-518

DDX5 Antibody (C-term) - Additional Information**Gene ID** 1655**Other Names**

Probable ATP-dependent RNA helicase DDX5, DEAD box protein 5, RNA helicase p68, DDX5, G17P1, HELR, HLR1

Target/Specificity

This DDX5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 491-518 amino acids from the C-terminal region of human DDX5.

Dilution

WB~~1:1000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DDX5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DDX5 Antibody (C-term) - Protein Information**Name** DDX5

Synonyms G17P1, HELR, HLR1

Function Involved in the alternative regulation of pre-mRNA splicing; its RNA helicase activity is necessary for increasing tau exon 10 inclusion and occurs in a RBM4-dependent manner. Binds to the tau pre-mRNA in the stem-loop region downstream of exon 10. The rate of ATP hydrolysis is highly stimulated by single-stranded RNA. Involved in transcriptional regulation; the function is independent of the RNA helicase activity. Transcriptional coactivator for androgen receptor AR but probably not ESR1. Synergizes with DDX17 and SRA1 RNA to activate MYOD1 transcriptional activity and involved in skeletal muscle differentiation. Transcriptional coactivator for p53/TP53 and involved in p53/TP53 transcriptional response to DNA damage and p53/TP53-dependent apoptosis. Transcriptional coactivator for RUNX2 and involved in regulation of osteoblast differentiation. Acts as a transcriptional repressor in a promoter-specific manner; the function probably involves association with histone deacetylases, such as HDAC1. As component of a large PER complex is involved in the inhibition of 3' transcriptional termination of circadian target genes such as PER1 and NR1D1 and the control of the circadian rhythms.


Cellular Location

Nucleus, Nucleus, nucleolus Nucleus speckle. Cytoplasm. Note=During the G0 phase, predominantly located in the nucleus. Cytoplasmic levels increase during the G1/S phase. During the M phase, located at the vicinity of the condensed chromosomes. At G1, localizes in the cytoplasm


DDX5 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DDX5 Antibody (C-term) - Images

Western blot analysis of DDX5 antibody (C-term) (Cat.#AP7459b) in T47D cell line lysates (35ug/lane). DDX5 (arrow) was detected using the purified Pab .



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with DDX5 antibody (C-term) (Cat. #AP7459b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

DDX5 Antibody (C-term) - Background

DDX5 is putative RNA helicases. The protein is implicated in a number of cellular processes involving alteration of RNA secondary structure, such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This protein is a RNA-dependent ATPase, and also a proliferation-associated nuclear antigen, specifically reacting with the simian virus 40 tumor antigen.

DDX5 Antibody (C-term) - References

Ong S.E., Mittler G. Nat. Methods 1:119-126(2004)
Daub H., Olsen J.V. Mol. Cell 31:438-448(2008)