

AXL Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7602d

Specification

AXL Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Isotype

WB, IHC-P, FC,E
P30530
Human
Rabbit
Polyclonal
Rabbit IgG
98337

AXL Antibody - Additional Information

Gene ID 558

Calculated MW

Other Names

Tyrosine-protein kinase receptor UFO, AXL oncogene, AXL, UFO

Target/Specificity

This AXL antibody is generated from rabbits immunized with a his tag recombinant protein of human AXL.

Dilution

WB~~1:2000 IHC-P~~1:50~100 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

AXL Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

AXL Antibody - Protein Information

Name AXL

Synonyms UFO

Function Receptor tyrosine kinase that transduces signals from the extracellular matrix into the



cytoplasm by binding growth factor GAS6 and which is thus regulating many physiological processes including cell survival, cell proliferation, migration and differentiation. Ligand binding at the cell surface induces dimerization and autophosphorylation of AXL. Following activation by ligand, AXL binds and induces tyrosine phosphorylation of PI3-kinase subunits PIK3R1, PIK3R2 and PIK3R3; but also GRB2, PLCG1, LCK and PTPN11. Other downstream substrate candidates for AXL are CBL, NCK2, SOCS1 and TNS2. Recruitment of GRB2 and phosphatidylinositol 3 kinase regulatory subunits by AXL leads to the downstream activation of the AKT kinase. GAS6/AXL signaling plays a role in various processes such as endothelial cell survival during acidification by preventing apoptosis, optimal cytokine signaling during human natural killer cell development, hepatic regeneration, gonadotropin-releasing hormone neuron survival and migration, platelet activation, or regulation of thrombotic responses. Also plays an important role in inhibition of Toll-like receptors (TLRs)-mediated innate immune response.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location


Highly expressed in metastatic colon tumors. Expressed in primary colon tumors. Weakly expressed in normal colon tissue.

AXL Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture


AXL Antibody - Images




All lanes: Anti-AXL Antibody at 1:2000 dilution Lane 1: Hela whole cell lysate Lane 2: DU145 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),



Peroxidase conjugated at 1/10000 dilution. Predicted band size : 98 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human colon carcinoma with AXL Antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of WiDr cells using AXL Antibody(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

AXL Antibody - Background

AXL, a member of the AXL/UFO subfamily of Tyr protein kinases, may function as a signal transducer between specific cell types of mesodermal origin. This Type I membrane protein has transforming potential in patients with chronic myeloproliferative disorder or chronic myelocytic leukemia. The protein contains 2 putative fibronectin type III domains and 2 putative immunoglobulin-like C2-type domains.

AXL Antibody - References

Lee, S.T., et al., Oncogene 8(12):3403-3410 (1993). Janssen, J.W., et al., Oncogene 6(11):2113-2120 (1991). O'Bryan, J.P., et al., Mol. Cell. Biol. 11(10):5016-5031 (1991). Partanen, J., et al., Proc. Natl. Acad. Sci. U.S.A. 87(22):8913-8917 (1990).