10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SRMS Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7719a

Specification

SRMS Antibody (N-term) - Product Information

Application	**WB, IHC-P,E**
Primary Accession	Q9H3Y6
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**54507**
Antigen Region	**19-48**

SRMS Antibody (N-term) - Additional Information

Gene ID 6725

Other Names
Tyrosine-protein kinase Srms, SRMS, C20orf148

Target/Specificity
This SRMS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 19-48 amino acids from the N-terminal region of human SRMS.

Dilution
WB~~1:1000
IHC-P~~1:50~100

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
SRMS Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SRMS Antibody (N-term) - Protein Information

Name SRMS

Synonyms C20orf148

Function Non-receptor tyrosine-protein kinase which phosphorylates DOK1 on tyrosine residues (PubMed:23822091). Also phosphorylates KHDRBS1/SAM68 and VIM on tyrosine residues (PubMed:29496907). Phosphorylation of KHDRBS1 is EGF-dependent (PubMed:29496907). Phosphorylates OTUB1, promoting deubiquitination of RPTOR (PubMed:35927303).

Cellular Location
Cytoplasm. Note=Localizes to punctate cytoplasmic structures.


Tissue Location
Highly expressed in most breast cancers (at protein level)

SRMS Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

SRMS Antibody (N-term) - Images



Western blot analysis of anti-SRMS Antibody (N-term) (Cat.#AP7719a) in HL60 cell line lysates (35ug/lane). SRMS(arrow) was detected using the purified Pab.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

SRMS Antibody (N-term) - Background

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the g phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The STE group (homologs of yeast Sterile 7, 11, 20 kinases) consists of 50 kinases related to the mitogen-activated protein kinase (MAPK) cascade families (Ste7/MAP2K, Ste11/MAP3K, and Ste20/MAP4K). MAP kinase cascades, consisting of a MAPK and one or more upstream regulatory kinases (MAPKKs) have been best characterized in the yeast pheromone response pathway. Pheromones bind to Ste cell surface receptors and activate yeast MAPK pathway.

SRMS Antibody (N-term) - References

Kohmura, N., et al., Mol. Cell. Biol. 14(10):6915-6925 (1994).