

**NGFB Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP7771c****Specification**

---

**NGFB Antibody (Center) - Product Information**

Application	IF, WB, IHC-P, FC,E
Primary Accession	<a href="#">P01138</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	26959
Antigen Region	84-115

**NGFB Antibody (Center) - Additional Information****Gene ID** 4803**Other Names**

Beta-nerve growth factor, Beta-NGF, NGF, NGFB

**Target/Specificity**

This NGFB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 84-115 amino acids from the Central region of human NGFB.

**Dilution**

IF~~1:10~50  
WB~~1:1000  
IHC-P~~1:10~50  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

NGFB Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**NGFB Antibody (Center) - Protein Information****Name** NGF

## Synonyms NGFB

**Function** Nerve growth factor is important for the development and maintenance of the sympathetic and sensory nervous systems (PubMed:[14976160](#), PubMed:[20978020](#)). Extracellular ligand for the NTRK1 and NGFR receptors, activates cellular signaling cascades to regulate neuronal proliferation, differentiation and survival (PubMed:[20978020](#)) (Probable). The immature NGF precursor (proNGF) functions as a ligand for the heterodimeric receptor formed by SORCS2 and NGFR, and activates cellular signaling cascades that lead to inactivation of RAC1 and/or RAC2, reorganization of the actin cytoskeleton and neuronal growth cone collapse. In contrast to mature NGF, the precursor form (proNGF) promotes neuronal apoptosis (in vitro) (By similarity). Inhibits metalloproteinase-dependent proteolysis of platelet glycoprotein VI (PubMed:[20164177](#)). Binds lysophosphatidylinositol and lysophosphatidylserine between the two chains of the homodimer. The lipid-bound form promotes histamine release from mast cells, contrary to the lipid-free form (By similarity).

## Cellular Location


Secreted. Endosome lumen {ECO:0000250|UniProtKB:P01139}. Note=ProNGF is endocytosed after binding to the cell surface receptor formed by SORT1 and NGFR {ECO:0000250|UniProtKB:P01139}

## NGFB Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## NGFB Antibody (Center) - Images




Confocal immunofluorescent analysis of NGFB Antibody (Center)(Cat#AP7771c) with MDA-MB231 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green).DAPI was used to stain


the cell nuclear (blue).



Western blot analysis of NGFB Antibody (Center) (Cat. #AP7771c) in NCI-H460 cell line lysates (35ug/lane). NGFB (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with NGFB antibody (Center) (Cat.#AP7771c), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



NGFB Antibody (Center) (Cat. #AP7771c) flow cytometric analysis of NCI-H460 cells (right

histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### **NGFB Antibody (Center) - Background**

NGFB is a member of the NGF-beta family. It is a secreted protein which homodimerizes and is incorporated into a larger complex. This protein has nerve growth stimulating activity and the complex is involved in the regulation of growth and the differentiation of sympathetic and certain sensory neurons. Mutations in the gene encoding NGFB have been associated with hereditary sensory and autonomic neuropathy, type 5 (HSAN5), and dysregulation of the gene's expression is associated with allergic rhinitis.

#### **NGFB Antibody (Center) - References**

Lang,U.E., Pharmacopsychiatry 41 (5), 196-199 (2008)  
Truzzi,F., J. Invest. Dermatol. 128 (8), 2031-2040 (2008)  
MacGrogan,D., J. Neurochem. 59 (4), 1381-1391 (1992)