

**CTGF Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP7778C****Specification**

---

**CTGF Antibody (Center) - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">P29279</a>
Other Accession	<a href="#">O9R1E9</a> , <a href="#">O19113</a> , <a href="#">P29268</a> , <a href="#">O18739</a> , <a href="#">NP_001892</a>
Reactivity	Human, Mouse
Predicted	Bovine, Pig, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	134-159

**CTGF Antibody (Center) - Additional Information****Gene ID** 1490**Other Names**

Connective tissue growth factor, CCN family member 2, Hypertrophic chondrocyte-specific protein 24, Insulin-like growth factor-binding protein 8, IBP-8, IGF-binding protein 8, IGFBP-8, CTGF, CCN2, HCS24, IGFBP8

**Target/Specificity**

This CTGF antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 134-159 amino acids from the Central region of human CTGF.

**Dilution**

WB~~1:1000  
IHC-P~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CTGF Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**CTGF Antibody (Center) - Protein Information****Name** CCN2 ([HGNC:2500](#))

**Function** Major connective tissue mitogen secreted by vascular endothelial cells. Promotes proliferation and differentiation of chondrocytes. Mediates heparin- and divalent cation-dependent cell adhesion in many cell types including fibroblasts, myofibroblasts, endothelial and epithelial cells. Enhances fibroblast growth factor- induced DNA synthesis.

**Cellular Location**

Secreted, extracellular space, extracellular matrix {ECO:0000250|UniProtKB:P29268}. Secreted {ECO:0000250|UniProtKB:P29268}

**Tissue Location**


Expressed in bone marrow and thymic cells. Also expressed one of two Wilms tumors tested.

**CTGF Antibody (Center) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CTGF Antibody (Center) - Images**



CTGF Antibody (Center) (Cat.#AP7778c) western blot analysis in mouse bladder tissue lysates (35ug/lane). This demonstrates the CTGF antibody detected the CTGF protein (arrow).



CTGF Antibody (Center) (Cat. #AP7778c) immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of CTGF Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

#### **CTGF Antibody (Center) - Background**

Connective tissue growth factor (CTGF), a member of the CCN family of secreted matricellular proteins, regulates fibrosis, angiogenesis, cell proliferation, apoptosis, tumor growth, and metastasis.

#### **CTGF Antibody (Center) - References**

Li, M.H., Mol. Cancer Res. 6 (10), 1649-1656 (2008)  
Bogatkevich, G.S., Am. J. Physiol. Lung Cell Mol. Physiol. 295 (4), L603-L611 (2008)  
Zhang, P., Tohoku J. Exp. Med. 215 (3), 199-206 (2008)