

IRAK3 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7804a**Specification**

IRAK3 Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O9Y616
Other Accession	NP_009130
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	45-77

IRAK3 Antibody (N-term) - Additional Information**Gene ID** 11213**Other Names**

Interleukin-1 receptor-associated kinase 3, IRAK-3, IL-1 receptor-associated kinase M, IRAK-M, IRAK3 {ECO:0000312|EMBL:AAH578001}

Target/Specificity

This IRAK3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 45-77 amino acids from the N-terminal region of human IRAK3.

Dilution

WB~~1:500

IHC-P~~1:100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IRAK3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

IRAK3 Antibody (N-term) - Protein Information**Name** IRAK3 {ECO:0000312|EMBL:AAH57800.1}**Function** Putative inactive protein kinase which regulates signaling downstream of immune

receptors including IL1R and Toll-like receptors (PubMed:[10383454](#), PubMed:[29686383](#)). Inhibits dissociation of IRAK1 and IRAK4 from the Toll-like receptor signaling complex by either inhibiting the phosphorylation of IRAK1 and IRAK4 or stabilizing the receptor complex (By similarity). Upon IL33-induced lung inflammation, positively regulates expression of IL6, CSF3, CXCL2 and CCL5 mRNAs in dendritic cells (PubMed:[29686383](#)).

Cellular Location

Cytoplasm. Nucleus. Note=In dendritic cells, translocates into the nucleus upon IL33 stimulation. {ECO:0000250|UniProtKB:Q8K4B2}

Tissue Location


Expressed in eosinophils, dendritic cells and/or monocytes (at protein level) (PubMed:29686383). Expressed predominantly in peripheral blood lymphocytes (PubMed:10383454)

IRAK3 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IRAK3 Antibody (N-term) - Images



All lanes : Anti-IRAK3 Antibody (N-term) at 1:500 dilution Lane 1: Raji whole cell lysate Lane 2: K562 whole cell lysate Lane 3: Mouse spleen tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 68 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of AP7804A on paraffin-embedded Human tonsil tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9.0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

IRAK3 Antibody (N-term) - Background

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the γ phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The tyrosine-like kinase (TLK) group consists of 40 tyrosine and serine-threonine kinases such as MLK (mixed-lineage kinase), LISK (LIMK/TESK), IRAK (interleukin-1 receptor-associated kinase), Raf, RIPK (receptor-interacting protein kinase), and STRK (activin and TGF-beta receptors) families.

IRAK3 Antibody (N-term) - References

Rosati, O., et al., Biochem. Biophys. Res. Commun. 293(5):1472-1477 (2002).
Wesche, H., et al., J. Biol. Chem. 274(27):19403-19410 (1999).