

YARS2 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7838b**Specification**

YARS2 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q9Y2Z4
Other Accession	Q5I0L3 , Q8BYL4
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	399-427

YARS2 Antibody (C-term) - Additional Information**Gene ID** 51067**Other Names**

Tyrosine--tRNA ligase, mitochondrial, Tyrosyl-tRNA synthetase, TyrRS, YARS2

Target/Specificity

This YARS2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 399-427 amino acids from the C-terminal region of human YARS2.

Dilution

WB~~1:1000

IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

YARS2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

YARS2 Antibody (C-term) - Protein Information**Name** YARS2**Function** Catalyzes the attachment of tyrosine to tRNA(Tyr) in a two- step reaction: tyrosine is

first activated by ATP to form Tyr-AMP and then transferred to the acceptor end of tRNA(Tyr).

Cellular Location


Mitochondrion matrix

YARS2 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

YARS2 Antibody (C-term) - Images



All lanes : Anti-YARS2 Antibody (C-term) at 1:1000 dilution Lane 1: K562 whole cell lysate Lane 2: CCRF-CEM whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 53 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with YARS2 antibody (C-term) (Cat.#AP7838b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

YARS2 Antibody (C-term) - Background

YARS2 catalyzes the attachment of tyrosine to tRNA(Tyr) in a two-step reaction: tyrosine is first activated by ATP to form Tyr-AMP and then transferred to the acceptor end of tRNA(Tyr).

YARS2 Antibody (C-term) - References

Bonnefond,L., Structure 15 (11), 1505-1516 (2007)

Bonnefond,L., RNA 11 (5), 558-562 (2005)

Bonnefond,L., Biochemistry 44 (12), 4805-4816 (2005)

YARS2 Antibody (C-term) - Citations

- [Mutation of the mitochondrial tyrosyl-tRNA synthetase gene, YARS2, causes myopathy, lactic acidosis, and sideroblastic anemia--MLASA syndrome.](#)