

AAK1 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7861B

Specification

AAK1 Antibody (C-term) - Product Information

Application WB,E
Primary Accession Q2M2I8

Other Accession POC1X8, F1SPM8, Q3UHIO, F1MH24

Reactivity Human

Predicted Bovine, Mouse, Pig, Rat

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 103885
Antigen Region 790-819

AAK1 Antibody (C-term) - Additional Information

Gene ID 22848

Other Names

AP2-associated protein kinase 1, Adaptor-associated kinase 1, AAK1, KIAA1048

Target/Specificity

This AAK1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 790-819 amino acids from the C-terminal region of human AAK1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

AAK1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

AAK1 Antibody (C-term) - Protein Information

Name AAK1

Synonyms KIAA1048



Function Regulates clathrin-mediated endocytosis by phosphorylating the AP2M1/mu2 subunit of the adaptor protein complex 2 (AP-2) which ensures high affinity binding of AP-2 to cargo membrane proteins during the initial stages of endocytosis (PubMed:17494869, PubMed:11877457, PubMed:11877461, PubMed:12952931, PubMed:14617351, PubMed:25653444). Isoform 1 and isoform 2 display similar levels of kinase activity towards AP2M1 (PubMed:17494869). Preferentially, may phosphorylate substrates on threonine residues (PubMed:11877457, PubMed:18657069). Regulates phosphorylation of other AP-2 subunits as well as AP-2 localization and AP-2-mediated internalization of ligand complexes (PubMed:12952931). Phosphorylates NUMB and regulates its cellular localization, promoting NUMB localization to endosomes (PubMed:18657069). Binds to and stabilizes the activated form of NOTCH1, increases its localization in endosomes and regulates its transcriptional activity (PubMed:21464124).

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:F1MH24}; Peripheral membrane protein {ECO:0000250|UniProtKB:F1MH24}. Membrane, clathrin-coated pit. Presynapse {ECO:0000250|UniProtKB:P0C1X8}. Note=Active when found in clathrin- coated pits at the plasma membrane. In neuronal cells, enriched at presynaptic terminals. In non-neuronal cells, enriched at leading edge of migrating cells. {ECO:0000250|UniProtKB:P0C1X8}

Tissue Location


Detected in brain, heart and liver. Isoform 1 is the predominant isoform in brain.

AAK1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

AAK1 Antibody (C-term) - Images



Western blot analysis of anti-AAK1 Antibody (C-term) (Cat.#AP7861b) in HepG2 cell line lysates (35ug/lane). AAK1 (arrow) was detected using the purified Pab.



Tel: 858.875.1900 Fax: 858.875.1999

AAK1 Antibody (C-term) - Background

Adaptor-related protein complex 2 (AP-2 complexes) functions during receptor-mediated endocytosis to trigger clathrin assembly, interact with membrane-bound receptors, and recruit encodytic accessory factors. AAK1 is a member of the SNF1 subfamily of Ser/Thr protein kinases. The protein interacts with and phosphorylates a subunit of the AP-2 complex, which promotes binding of AP-2 to sorting signals found in membrane-bound receptors and subsequent receptor endocytosis. Its kinase activity is stimulated by clathrin.

AAK1 Antibody (C-term) - References

Henderson, D.M., Mol. Biol. Cell 18 (7), 2698-2706 (2007) Takahashi, T., Cancer Res. 66 (24), 11932-11937 (2006) Schmid, E.M., PLoS Biol. 4 (9), E262 (2006)