10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MEK3 (MAP2K3) Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7915a

Specification

MEK3 (MAP2K3) Antibody (N-term) - Product Information

Application	**WB, IHC-P,E**
Primary Accession	P46734
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Antigen Region	**15-45**

MEK3 (MAP2K3) Antibody (N-term) - Additional Information

Gene ID 5606

Other Names
Dual specificity mitogen-activated protein kinase kinase 3, MAP kinase kinase 3, MAPKK 3, MAPK/ERK kinase 3, MEK 3, Stress-activated protein kinase kinase 2, SAPK kinase 2, SAPKK-2, SAPKK2, MAP2K3, MEK3, MKK3, PRKMK3, SKK2

Target/Specificity
This MEK3 (MAP2K3) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 15-45 amino acids from the N-terminal region of human MEK3 (MAP2K3).

Dilution
WB~~1:1000
IHC-P~~1:50~100

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
MEK3 (MAP2K3) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MEK3 (MAP2K3) Antibody (N-term) - Protein Information

Name MAP2K3

Synonyms MEK3, MKK3, PRKMK3, SKK2

Function Dual specificity kinase. Is activated by cytokines and environmental stress in vivo. Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in the MAP kinase p38. Part of a signaling cascade that begins with the activation of the adrenergic receptor ADRA1B and leads to the activation of MAPK14.


Tissue Location
Abundant expression is seen in the skeletal muscle. It is also widely expressed in other tissues

MEK3 (MAP2K3) Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

MEK3 (MAP2K3) Antibody (N-term) - Images




Western blot analysis of anti-MKK3 Pab (Cat. #AP7915a) in Jurkat cell lysate. MKK3 (arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



IHC of matched glioma core (top row) and rim (middle row) samples as well as normal brain control (bottom row) from a glioma invasion specific tissue microarray. Insets, individual tumor cells (black squares) are presented at higher magnification to facilitate interpretation of the staining pattern. Staining for MKK3 and pMKK3 exhibits strong signal in majority of invasive cells (B) compared with stationary cells from tumor core (A). Normal brain control shows only weak staining in reactive astrocytes (C).

MEK3 (MAP2K3) Antibody (N-term) - Background

MAP2K3 is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is activated by mitogenic and environmental stress, and participates in the MAP kinase-mediated signaling cascade. It phosphorylates and thus activates MAPK14/p38-MAPK. This kinase can be activated by insulin, and is necessary for the expression of glucose transporter. Expression of RAS oncogene is found to result in the accumulation of the active form of this kinase, which thus leads to the constitutive activation of MAPK14, and confers oncogenic transformation of primary cells. The inhibition of this kinase is involved in the pathogenesis of Yersina pseudotuberculosis.

MEK3 (MAP2K3) Antibody (N-term) - References

Yustein, J.T., et al., Oncogene 22(40):6129-6141 (2003).
Edlund, S., et al., Mol. Biol. Cell 14(2):529-544 (2003).
Lim, S., et al., J. Biol. Chem. 277(28):25040-25046 (2002).
Han, Q., et al., J. Biol. Chem. 277(50):48379-48385 (2002).
Wang, W., et al., Mol. Cell. Biol. 22(10):3389-3403 (2002).

MEK3 (MAP2K3) Antibody (N-term) - Citations

- MAP-ing glioma invasion: mitogen-activated protein kinase kinase 3 and p38 drive glioma invasion and progression and predict patient survival.