

**BRD4 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8051b****Specification**

---

**BRD4 Antibody (C-term) - Product Information**

Application	IHC-P,E
Primary Accession	<a href="#">O60885</a>
Other Accession	<a href="#">Q9ESU6</a>
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	1313-1342

**BRD4 Antibody (C-term) - Additional Information****Gene ID** 23476**Other Names**

Bromodomain-containing protein 4, Protein HUNK1, BRD4, HUNK1

**Target/Specificity**

This BRD4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1313~1342 amino acids from the C-terminal region of human BRD4.

**Dilution**

IHC-P~~1:50~100

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

BRD4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**BRD4 Antibody (C-term) - Protein Information****Name** BRD4**Synonyms** HUNK1

**Function** Chromatin reader protein that recognizes and binds acetylated histones and plays a key role in transmission of epigenetic memory across cell divisions and transcription regulation (PubMed:[23086925](#), PubMed:[23317504](#), PubMed:[20871596](#), PubMed:[29176719](#)). Remains associated with acetylated chromatin throughout the entire cell cycle and provides epigenetic memory for postmitotic G1 gene transcription by preserving acetylated chromatin status and maintaining high-order chromatin structure (PubMed:[23589332](#), PubMed:[23317504](#), PubMed:[22334664](#)). During interphase, plays a key role in regulating the transcription of signal-inducible genes by associating with the P-TEFb complex and recruiting it to promoters (PubMed:[23589332](#), PubMed:[19596240](#), PubMed:[16109377](#), PubMed:[16109376](#), PubMed:[24360279](#)). Also recruits P-TEFb complex to distal enhancers, so called anti-pause enhancers in collaboration with JMJD6 (PubMed:[23589332](#), PubMed:[19596240](#), PubMed:[16109377](#), PubMed:[16109376](#), PubMed:[24360279](#)). BRD4 and JMJD6 are required to form the transcriptionally active P-TEFb complex by displacing negative regulators such as HEXIM1 and 7SKsnRNA complex from P-TEFb, thereby transforming it into an active form that can then phosphorylate the C-terminal domain (CTD) of RNA polymerase II (PubMed:[23589332](#), PubMed:[19596240](#), PubMed:[16109377](#), PubMed:[16109376](#), PubMed:[24360279](#)). Regulates differentiation of naive CD4(+) T-cells into T-helper Th17 by promoting recruitment of P-TEFb to promoters (By similarity). Promotes phosphorylation of 'Ser-2' of the C-terminal domain (CTD) of RNA polymerase II (PubMed:[23086925](#)). According to a report, directly acts as an atypical protein kinase and mediates phosphorylation of 'Ser-2' of the C-terminal domain (CTD) of RNA polymerase II; these data however need additional evidences in vivo (PubMed:[22509028](#)). In addition to acetylated histones, also recognizes and binds acetylated RELA, leading to further recruitment of the P-TEFb complex and subsequent activation of NF-kappa-B (PubMed:[19103749](#)). Also acts as a regulator of p53/TP53-mediated transcription: following phosphorylation by CK2, recruited to p53/TP53 specific target promoters (PubMed:[23317504](#)).

#### **Cellular Location**

Nucleus. Chromosome. Note=Associates with acetylated chromatin (PubMed:21890894, PubMed:16109376). Released from chromatin upon deacetylation of histones that can be triggered by different signals such as activation of the JNK pathway or nocodazole treatment (PubMed:21890894, PubMed:16109376). Preferentially localizes to mitotic chromosomes, while it does not localize to meiotic chromosomes (PubMed:21890894, PubMed:16109376).

#### **Tissue Location**

Ubiquitously expressed.


#### **BRD4 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **BRD4 Antibody (C-term) - Images**





Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



Subnuclear distribution of cellular proteins. CHO Bgl40 cells grown on coverslips were either directly or after treatment with 0.5% Triton X-100, incubated with antibodies against Brd4 (images 4 and 5). PI, propidium iodide staining of cellular DNA (images 6). WC, whole cells.

#### BRD4 Antibody (C-term) - Background

BRD4 is homologous to the murine protein MCAP, which associates with chromosomes during mitosis, and to the human RING3 protein, a serine/threonine kinase. Each of these proteins contains two bromodomains, a conserved sequence motif which may be involved in chromatin targeting. The gene has been implicated as the chromosome 19 target of translocation t(15;19)(q13;p13.1), which defines an upper respiratory tract carcinoma in young people.

#### BRD4 Antibody (C-term) - References

Maruyama, T., et al., Mol. Cell. Biol. 22(18):6509-6520 (2002).  
French, C.A., et al., Am. J. Pathol. 159(6):1987-1992 (2001).  
Dey, A., et al., Mol. Cell. Biol. 20(17):6537-6549 (2000).

#### BRD4 Antibody (C-term) - Citations

- [BRD4 regulates fructose-inducible lipid accumulation-related genes in the mouse liver.](#)
- [Amino acid substitutions that specifically impair the transcriptional activity of papillomavirus E2 affect binding to the long isoform of Brd4.](#)
- [Characterization of the functional activities of the bovine papillomavirus type 1 E2 protein single-chain heterodimers.](#)
- [Association of bovine papillomavirus E2 protein with nuclear structures in vivo.](#)