

GSK3 beta (GSK3B) Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8121a**Specification**

GSK3 beta (GSK3B) Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	P49841
Other Accession	P18266 , Q9WV60
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	4-34

GSK3 beta (GSK3B) Antibody (N-term) - Additional Information**Gene ID** 2932**Other Names**

Glycogen synthase kinase-3 beta, GSK-3 beta, Serine/threonine-protein kinase GSK3B, GSK3B

Target/Specificity

This GSK3 beta (GSK3B) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 4-34 amino acids from the N-terminal region of human GSK3 beta (GSK3B).

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GSK3 beta (GSK3B) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

GSK3 beta (GSK3B) Antibody (N-term) - Protein Information**Name** GSK3B ([HGNC:4617](#))

Function Constitutively active protein kinase that acts as a negative regulator in the hormonal control of glucose homeostasis, Wnt signaling and regulation of transcription factors and microtubules, by phosphorylating and inactivating glycogen synthase (GYS1 or GYS2), EIF2B, CTNNB1/beta-catenin, APC, AXIN1, DPYSL2/CRMP2, JUN, NFATC1/NFATC, MAPT/TAU and MACF1 (PubMed:[1846781](#), PubMed:[9072970](#), PubMed:[14690523](#), PubMed:[20937854](#), PubMed:[12554650](#), PubMed:[11430833](#), PubMed:[16484495](#)). Requires primed phosphorylation of the majority of its substrates (PubMed:[11430833](#), PubMed:[16484495](#)). In skeletal muscle, contributes to insulin regulation of glycogen synthesis by phosphorylating and inhibiting GYS1 activity and hence glycogen synthesis (PubMed:[8397507](#)). May also mediate the development of insulin resistance by regulating activation of transcription factors (PubMed:[8397507](#)). Regulates protein synthesis by controlling the activity of initiation factor 2B (EIF2BE/EIF2B5) in the same manner as glycogen synthase (PubMed:[8397507](#)). In Wnt signaling, GSK3B forms a multimeric complex with APC, AXIN1 and CTNNB1/beta-catenin and phosphorylates the N-terminus of CTNNB1 leading to its degradation mediated by ubiquitin/proteasomes (PubMed:[12554650](#)). Phosphorylates JUN at sites proximal to its DNA-binding domain, thereby reducing its affinity for DNA (PubMed:[1846781](#)). Phosphorylates NFATC1/NFATC on conserved serine residues promoting NFATC1/NFATC nuclear export, shutting off NFATC1/NFATC gene regulation, and thereby opposing the action of calcineurin (PubMed:[9072970](#)). Phosphorylates MAPT/TAU on 'Thr-548', decreasing significantly MAPT/TAU ability to bind and stabilize microtubules (PubMed:[14690523](#)). MAPT/TAU is the principal component of neurofibrillary tangles in Alzheimer disease (PubMed:[14690523](#)). Plays an important role in ERBB2-dependent stabilization of microtubules at the cell cortex (PubMed:[20937854](#)). Phosphorylates MACF1, inhibiting its binding to microtubules which is critical for its role in bulge stem cell migration and skin wound repair (By similarity). Probably regulates NF-kappa-B (NFKB1) at the transcriptional level and is required for the NF-kappa-B-mediated anti-apoptotic response to TNF-alpha (TNF/TNFA) (By similarity). Negatively regulates replication in pancreatic beta-cells, resulting in apoptosis, loss of beta-cells and diabetes (By similarity). Through phosphorylation of the anti-apoptotic protein MCL1, may control cell apoptosis in response to growth factors deprivation (By similarity). Phosphorylates MUC1 in breast cancer cells, decreasing the interaction of MUC1 with CTNNB1/beta-catenin (PubMed:[9819408](#)). Is necessary for the establishment of neuronal polarity and axon outgrowth (PubMed:[20067585](#)). Phosphorylates MARK2, leading to inhibition of its activity (By similarity). Phosphorylates SIK1 at 'Thr-182', leading to sustainment of its activity (PubMed:[18348280](#)). Phosphorylates ZC3HAV1 which enhances its antiviral activity (PubMed:[22514281](#)). Phosphorylates SNAI1, leading to its BTRC-triggered ubiquitination and proteasomal degradation (PubMed:[15448698](#), PubMed:[15647282](#)). Phosphorylates SFPQ at 'Thr-687' upon T-cell activation (PubMed:[20932480](#)). Phosphorylates NR1D1 at 'Ser-55' and 'Ser-59' and stabilizes it by protecting it from proteasomal degradation. Regulates the circadian clock via phosphorylation of the major clock components including BMAL1, CLOCK and PER2 (PubMed:[19946213](#), PubMed:[28903391](#)). Phosphorylates FBXL2 at 'Thr-404' and primes it for ubiquitination by the SCF(FBXO3) complex and proteasomal degradation (By similarity). Phosphorylates CLOCK at 'Ser-427' and targets it for proteasomal degradation (PubMed:[19946213](#)). Phosphorylates BMAL1 at 'Ser-17' and 'Ser-21' and primes it for ubiquitination and proteasomal degradation (PubMed:[28903391](#)). Phosphorylates OGT at 'Ser-3' or 'Ser-4' which positively regulates its activity. Phosphorylates MYCN in neuroblastoma cells which may promote its degradation (PubMed:[24391509](#)). Regulates the circadian rhythmicity of hippocampal long-term potentiation and BMAL1 and PER2 expression (By similarity). Acts as a regulator of autophagy by mediating phosphorylation of KAT5/TIP60 under starvation conditions, activating KAT5/TIP60 acetyltransferase activity and promoting acetylation of key autophagy regulators, such as ULK1 and RUBCNL/Pacer (PubMed:[30704899](#)). Negatively regulates extrinsic apoptotic signaling pathway via death domain receptors. Promotes the formation of an anti-apoptotic complex, made of DDX3X, BRIC2 and GSK3B, at death receptors, including TNFRSF10B. The anti-apoptotic function is most effective with weak apoptotic signals and can be overcome by stronger stimulation (PubMed:[18846110](#)). Phosphorylates E2F1, promoting the interaction between E2F1 and USP11, stabilizing E2F1 and promoting its activity (PubMed:[17050006](#), PubMed:[28992046](#)). Phosphorylates mTORC2 complex component RICTOR at 'Thr-1695' which facilitates FBXW7-mediated ubiquitination and subsequent degradation of RICTOR (PubMed:[25897075](#)). Phosphorylates FXR1, promoting FXR1 ubiquitination by the SCF(FBXO4) complex and FXR1 degradation by the proteasome (By similarity). Phosphorylates

interleukin-22 receptor subunit IL22RA1, preventing its proteasomal degradation (By similarity).

Cellular Location

Cytoplasm. Nucleus. Cell membrane. Note=The phosphorylated form shows localization to cytoplasm and cell membrane (PubMed:20937854). The MEMO1-RHOA-DIAPH1 signaling pathway controls localization of the phosphorylated form to the cell membrane (PubMed:20937854)

Tissue Location


Expressed in testis, thymus, prostate and ovary and weakly expressed in lung, brain and kidney. Colocalizes with EIF2AK2/PKR and TAU in the Alzheimer disease (AD) brain

GSK3 beta (GSK3B) Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GSK3 beta (GSK3B) Antibody (N-term) - Images



The anti-GSK3B Pab (Cat. #AP8121a) is used in Western blot to detect GSK3B in 293 cell lysate.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

GSK3 beta (GSK3B) Antibody (N-term) - Background

GSK3B, a member of the Ser/Thr protein kinase family, participates in the Wnt signaling pathway. It is implicated in the hormonal control of several regulatory proteins including glycogen synthase, MYB and the transcription factor JUN. GSK3B phosphorylates JUN at sites proximal to its DNA-binding domain, thereby reducing its affinity for DNA. The protein is inhibited when phosphorylated by AKT1 or ILK1. Expressed is detected in testis, thymus, prostate and ovary, and weakly in lung, brain and kidney.

GSK3 beta (GSK3B) Antibody (N-term) - References

Strausberg, R.L., et al., Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903 (2002).
Dajani, R., et al., Cell 105(6):721-732 (2001).
Lau, K.F., et al., Genomics 60(2):121-128 (1999).
Rhoads, A.R., et al., Mol. Psychiatry 4(5):437-442 (1999).
Delcommenne, M., et al., Proc. Natl. Acad. Sci. U.S.A. 95(19):11211-11216 (1998).

GSK3 beta (GSK3B) Antibody (N-term) - Citations

- [Axl mediates tumor invasion and chemosensitivity through PI3K/Akt signaling pathway and is transcriptionally regulated by slug in breast carcinoma.](#)