

AK3 Antibody (C-term H38)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8132b**Specification**

AK3 Antibody (C-term H38) - Product Information

Application	WB, IHC-P,E
Primary Accession	P27144
Other Accession	O9WUS0 , O9WUR9
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	25268
Antigen Region	195-223

AK3 Antibody (C-term H38) - Additional Information**Gene ID** 205**Other Names**

Adenylate kinase 4, mitochondrial {ECO:0000255|HAMAP-Rule:MF_03170}, AK 4 {ECO:0000255|HAMAP-Rule:MF_03170}, 27410 {ECO:0000255|HAMAP-Rule:MF_03170}, 2746 {ECO:0000255|HAMAP-Rule:MF_03170}, Adenylate kinase 3-like {ECO:0000255|HAMAP-Rule:MF_03170}, GTP:AMP phosphotransferase AK4 {ECO:0000255|HAMAP-Rule:MF_03170}, AK4 {ECO:0000255|HAMAP-Rule:MF_03170}

Target/Specificity

This AK3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 195-223 amino acids from the C-terminal region of human AK3.

Dilution

WB~~1:1000
IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

AK3 Antibody (C-term H38) is for research use only and not for use in diagnostic or therapeutic procedures.

AK3 Antibody (C-term H38) - Protein Information

Name AK4 {ECO:0000255|HAMAP-Rule:MF_03170}

Function Involved in maintaining the homeostasis of cellular nucleotides by catalyzing the interconversion of nucleoside phosphates (PubMed:[19073142](#), PubMed:[19766732](#), PubMed:[23416111](#), PubMed:[24767988](#)). Efficiently phosphorylates AMP and dAMP using ATP as phosphate donor, but phosphorylates only AMP when using GTP as phosphate donor (PubMed:[19073142](#), PubMed:[19766732](#), PubMed:[23416111](#)). Also displays broad nucleoside diphosphate kinase activity (PubMed:[19073142](#), PubMed:[19766732](#), PubMed:[23416111](#)). Plays a role in controlling cellular ATP levels by regulating phosphorylation and activation of the energy sensor protein kinase AMPK (PubMed:[24767988](#), PubMed:[26980435](#)). Plays a protective role in the cellular response to oxidative stress (PubMed:[19130895](#), PubMed:[23474458](#), PubMed:[26980435](#)).

Cellular Location

Mitochondrion matrix {ECO:0000255|HAMAP- Rule:MF_03170, ECO:0000269|PubMed:11485571, ECO:0000269|PubMed:19766732, ECO:0000269|PubMed:26980435}

Tissue Location


Highly expressed in kidney, moderately expressed in heart and liver and weakly expressed in brain

AK3 Antibody (C-term H38) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


AK3 Antibody (C-term H38) - Images



The anti-AK3 Pab (Cat. #AP8132b) is used in Western blot to detect AK3 in mouse kidney tissue lysate.



Western blot analysis of AK3 (arrow) using rabbit polyclonal AK3 Antibody (C-term H38) (Cat. #AP8132b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the AK3 gene (Lane 2) (Origene Technologies).



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

AK3 Antibody (C-term H38) - Background

AK3 is a member of the adenylate kinase family of enzymes. The encoded protein is localized to the mitochondrial matrix. Adenylate kinases regulate the adenine and guanine nucleotide compositions within a cell by catalyzing the reversible transfer of phosphate group among these nucleotides. Five isozymes of adenylate kinase have been identified in vertebrates. Expression of these isozymes is tissue-specific and developmentally regulated.

AK3 Antibody (C-term H38) - References

- Van Rompay, A.R., et al., Eur. J. Biochem. 261(2):509-517 (1999).
- Yoneda, T., et al., Brain Res. Mol. Brain Res. 62(2):187-195 (1998).
- Xu, G., et al., Genomics 13(3):537-542 (1992).
- Robson, E.B., et al., Cytogenet. Cell Genet. 32 (1-4), 144-152 (1982).