

PTP gamma Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP8414a

Specification

PTP gamma Antibody (Center) - Product Information

Application WB,E
Primary Accession P23470

Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 162003
Antigen Region 751-782

PTP gamma Antibody (Center) - Additional Information

Gene ID 5793

Other Names

Receptor-type tyrosine-protein phosphatase gamma, Protein-tyrosine phosphatase gamma, R-PTP-gamma, PTPRG, PTPG

Target/Specificity

This PTP gamma antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 751-782 amino acids from the Central region of human PTP gamma.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PTP gamma Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PTP gamma Antibody (Center) - Protein Information

Name PTPRG

Synonyms PTPG



Function Possesses tyrosine phosphatase activity.

Cellular Location

Membrane; Single-pass type I membrane protein

Tissue Location


Found in a variety of tissues.

PTP gamma Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

PTP gamma Antibody (Center) - Images



The anti-PTPgamma Center Pab (Cat. #AP8414a) is used in Western blot to detect PTPgamma in mouse brain tissue lysate.

PTP gamma Antibody (Center) - Background

PTPgamma is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP possesses an extracellular region, a single transmembrane region, and two tandem intracytoplasmic catalytic domains, and thus represents a receptor-type PTP. The extracellular region of this PTP contains a carbonic anhydrase-like (CAH) domain, which is also found in the extracellular region of PTPRBETA/ZETA. The gene is located in a chromosomal region that is frequently deleted in renal cell carcinoma and lung carcinoma, thus is thought to be a candidate tumor suppressor gene.

PTP gamma Antibody (Center) - References





Liu, S., et al., Breast Cancer Res. Treat. 71(1):21-35 (2002). Kastury, K., et al., Genomics 32(2):225-235 (1996).

Barnea, G., et al., Mol. Cell. Biol. 13(3):1497-1506 (1993).

LaForgia, S., et al., Proc. Natl. Acad. Sci. U.S.A. 88(11):5036-5040 (1991).

Krueger, N.X., et al., EMBO J. 9(10):3241-3252 (1990).