10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# HSPC142 Antibody (Center)

**Purified Rabbit Polyclonal Antibody (Pab)**

**Catalog # AP8573c**

## Specification

### HSPC142 Antibody (Center) - Product Information

| | |
|---|---|
| Application | **WB, IHC-P, FC,E** |
| Primary Accession | Q9NWV8 |
| Other Accession | Q5XIJ6, Q3UI43, Q08E57 |
| Reactivity | **Human** |
| Predicted | **Bovine, Mouse, Rat** |
| Host | **Rabbit** |
| Clonality | **Polyclonal** |
| Isotype | **Rabbit IgG** |
| Calculated MW | **36560** |
| Antigen Region | **116-143** |

### HSPC142 Antibody (Center) - Additional Information

**Gene ID** 29086

**Other Names**
BRISC and BRCA1-A complex member 1, Mediator of RAP80 interactions and targeting subunit of 40 kDa, New component of the BRCA1-A complex, BABAM1, C19orf62, MERIT40, NBA1

**Target/Specificity**
This HSPC142 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 116-143 amino acids from the Central region of human HSPC142.

**Dilution**
WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

**Format**
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**
HSPC142 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

### HSPC142 Antibody (Center) - Protein Information

**Name** BABAM1

**Function** Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'- linked ubiquitin on histones H2A and H2AX. In the BRCA1-A complex, it is required for the complex integrity and its localization at DSBs. Component of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin in various substrates (PubMed:24075985, PubMed:26195665). In these 2 complexes, it is probably required to maintain the stability of BABAM2 and help the 'Lys-63'-linked deubiquitinase activity mediated by BRCC3/BRCC36 component. The BRISC complex is required for normal mitotic spindle assembly and microtubule attachment to kinetochores via its role in deubiquitinating NUMA1 (PubMed:26195665). Plays a role in interferon signaling via its role in the deubiquitination of the interferon receptor IFNAR1; deubiquitination increases IFNAR1 activity by enhancing its stability and cell surface expression (PubMed:24075985). Down-regulates the response to bacterial lipopolysaccharide (LPS) via its role in IFNAR1 deubiquitination (PubMed:24075985).


**Cellular Location**
Cytoplasm. Nucleus Note=Localizes at sites of DNA damage at double-strand breaks (DSBs)

## HSPC142 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture


## HSPC142 Antibody (Center) - Images





Western blot analysis of HSPC142 Antibody (Center) (Cat.#AP8573c) in MDA-MB231 cell line

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

lysates (35ug/lane). HSPC142 (arrow) was detected using the purified Pab.




Formalin-fixed and paraffin-embedded human brain reacted with HSPC142 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.




Flow cytometric analysis of MDA-MB468 cells using HSPC142 Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram)FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

**HSPC142 Antibody (Center) - Background**

Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX. In the BRCA1-A complex, it is required for the complex integrity and its localization at DSBs. Probably also plays a role as a component of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin. In these 2 complexes, it is probably required to maintain the stability of BRE/BRCC45 and help the 'Lys-63'-linked deubiquitinase activity mediated by BRCC3/BRCC36. component.

**HSPC142 Antibody (Center) - References**

Shao,G., et.al., Genes Dev. 23 (6), 740-754 (2009)
Ewing,R.M., et.al., Mol. Syst. Biol. 3, 89 (2007)