

#### **KLC1 Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP8637c

#### Specification

## KLC1 Antibody (Center) - Product Information

| Application<br>Primary Accession<br>Other Accession<br>Reactivity<br>Predicted<br>Host<br>Clonality | WB, IHC-P, FC,E<br><u>007866</u><br><u>P37285</u><br>Human<br>Rat<br>Rabbit<br>Polyclonal |
|---|---|
| Clonality   | Polyclonal  |
| Isotype   | Rabbit IgG  |
| Antigen Region  | 389-415   |

#### **KLC1 Antibody (Center) - Additional Information**

Gene ID 3831

Other Names Kinesin light chain 1, KLC 1, KLC1, KLC, KNS2

**Target/Specificity** This KLC1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 389-415 amino acids from the Central region of human KLC1.

**Dilution** WB~~1:1000 IHC-P~~1:100 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

KLC1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## KLC1 Antibody (Center) - Protein Information

Name KLC1



### Synonyms KLC, KNS2

**Function** Kinesin is a microtubule-associated force-producing protein that may play a role in organelle transport (PubMed:<u>21385839</u>). The light chain may function in coupling of cargo to the heavy chain or in the modulation of its ATPase activity (By similarity).

**Cellular Location** Cell projection, growth cone {ECO:0000250|UniProtKB:P37285}. Cytoplasmic vesicle. Cytoplasm, cytoskeleton


**Tissue Location** Found in a variety of tissues. Mostly abundant in brain and spine.

#### KLC1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

#### KLC1 Antibody (Center) - Images




KLC1 Antibody (Center) (Cat.# AP8637c) western blot analysis in Hela cell line lysates (35ug/lane).This demonstrates the KLC1 antibody detected the KLC1 protein (arrow).






Formalin-fixed and paraffin-embedded human hepatocarcinoma with KLC1 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Formalin-fixed and paraffin-embedded H.brain, H.cerebellum tissue reacted with KLC1 Antibody (Center) (Cat#AP8637c).



Flow cytometric analysis of widr cells using KLC1 Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

# KLC1 Antibody (Center) - Background

Kinesin is a microtubule-associated force-producing protein that may play a role in organelle transport. The light chain may function in coupling of cargo to the heavy chain or in the modulation



of its ATPase activity.

# KLC1 Antibody (Center) - References

Chernajovsky,Y., et.al., DNA Cell Biol. 15 (11), 965-974 (1996) Gyoeva,F.K., et.al., J. Cell. Sci. 113 (PT 11), 2047-2054 (2000)

**KLC1 Antibody (Center) - Citations** 

• Alterations in axonal transport motor proteins in sporadic and experimental Parkinson\'s disease.