

FLCN Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8658c**Specification**

FLCN Antibody (Center) - Product Information

| | |
|-------------------|------------------------|
| Application | WB, IHC-P,E |
| Primary Accession | Q8NFG4 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 64473 |
| Antigen Region | 325-354 |

FLCN Antibody (Center) - Additional Information**Gene ID** 201163**Other Names**

Folliculin, BHD skin lesion fibrofolliculoma protein, Birt-Hogg-Dube syndrome protein, FLCN, BHD

Target/Specificity

This FLCN antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 325-354 amino acids from the Central region of human FLCN.

DilutionWB~~1:1000
IHC-P~~1:50~100**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FLCN Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

FLCN Antibody (Center) - Protein Information**Name** FLCN {ECO:0000303|PubMed:15657874, ECO:0000312|HGNC:HGNC:27310}**Function** Multi-functional protein, involved in both the cellular response to amino acid availability and in the regulation of glycolysis (PubMed:[17028174](#), PubMed:[18663353](#), PubMed:[21209915](#),

PubMed:[24081491](#), PubMed:[24095279](#), PubMed:[31704029](#), PubMed:[31672913](#), PubMed:[34381247](#), PubMed:[32612235](#), PubMed:[36103527](#), PubMed:[37079666](#)). GTPase-activating protein that plays a key role in the cellular response to amino acid availability through regulation of the non-canonical mTORC1 signaling cascade controlling the MiT/TFE factors TFE3 and TFE3 (PubMed:[17028174](#), PubMed:[18663353](#), PubMed:[21209915](#), PubMed:[24081491](#), PubMed:[24095279](#), PubMed:[24448649](#), PubMed:[31704029](#), PubMed:[31672913](#), PubMed:[32612235](#), PubMed:[36103527](#), PubMed:[37079666](#)). Activates mTORC1 by acting as a GTPase-activating protein: specifically stimulates GTP hydrolysis by RagC/RRAGC or RagD/RRAGD, promoting the conversion to the GDP-bound state of RagC/RRAGC or RagD/RRAGD, and thereby activating the kinase activity of mTORC1 (PubMed:[24095279](#), PubMed:[31704029](#), PubMed:[31672913](#), PubMed:[32612235](#), PubMed:[37079666](#)). The GTPase-activating activity is inhibited during starvation and activated in presence of nutrients (PubMed:[31672913](#), PubMed:[32612235](#)). Acts as a key component for non- canonical mTORC1-dependent control of the MiT/TFE factors TFE3 and TFE3, while it is not involved in mTORC1-dependent phosphorylation of canonical RPS6KB1/S6K1 and EIF4EBP1/4E-BP1 (PubMed:[21209915](#), PubMed:[24081491](#), PubMed:[31672913](#), PubMed:[32612235](#)). In low-amino acid conditions, the lysosomal folliculin complex (LFC) is formed on the membrane of lysosomes, which inhibits the GTPase-activating activity of FLCN, inactivates mTORC1 and maximizes nuclear translocation of TFE3 and TFE3 (PubMed:[31672913](#)). Upon amino acid restimulation, RagA/RRAGA (or RagB/RRAGB) nucleotide exchange promotes disassembly of the LFC complex and liberates the GTPase-activating activity of FLCN, leading to activation of mTORC1 and subsequent cytoplasmic retention of TFE3 and TFE3 (PubMed:[31672913](#)). Indirectly acts as a positive regulator of Wnt signaling by promoting mTOR-dependent cytoplasmic retention of MiT/TFE factor TFE3 (PubMed:[31272105](#)). Required for the exit of hematopoietic stem cell from pluripotency by promoting mTOR-dependent cytoplasmic retention of TFE3, thereby increasing Wnt signaling (PubMed:[30733432](#)). Acts as an inhibitor of browning of adipose tissue by regulating mTOR-dependent cytoplasmic retention of TFE3 (By similarity). Involved in the control of embryonic stem cells differentiation; together with LAMTOR1 it is necessary to recruit and activate RagC/RRAGC and RagD/RRAGD at the lysosomes, and to induce exit of embryonic stem cells from pluripotency via non-canonical, mTOR- independent TFE3 inactivation (By similarity). In response to flow stress, regulates STK11/LKB1 accumulation and mTORC1 activation through primary cilia: may act by recruiting STK11/LKB1 to primary cilia for activation of AMPK resided at basal bodies, causing mTORC1 down- regulation (PubMed:[27072130](#)). Together with FNIP1 and/or FNIP2, regulates autophagy: following phosphorylation by ULK1, interacts with GABARAP and promotes autophagy (PubMed:[25126726](#)). Required for starvation-induced perinuclear clustering of lysosomes by promoting association of RILP with its effector RAB34 (PubMed:[27113757](#)). Regulates glycolysis by binding to lactate dehydrogenase LDHA, acting as an uncompetitive inhibitor (PubMed:[34381247](#)).

Cellular Location

Lysosome membrane. Cytoplasm, cytosol. Cell projection, cilium. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Nucleus
Note=Localizes to lysosome membrane in amino acid-depleted conditions and relocates to the cytosol upon refeeding (PubMed:[24095279](#), PubMed:[29848618](#), PubMed:[31672913](#)). Colocalizes with FNIP1 and FNIP2 in the cytoplasm (PubMed:[17028174](#), PubMed:[18663353](#)). Also localizes to motile and non-motile cilia, centrosomes and the mitotic spindle (PubMed:[23784378](#)).

Tissue Location

Expressed in most tissues tested, including skin, lung, kidney, heart, testis and stomach.


FLCN Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FLCN Antibody (Center) - Images



Western blot analysis of FLCN Antibody (Center) (Cat. #AP8658c) in CEM cell line lysates (35ug/lane). FLCN (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human lung carcinoma reacted with FLCN Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

FLCN Antibody (Center) - Background

FLCN may play a role in the pathogenesis of an uncommon form of kidney cancer through its association with an inherited disorder of the hair follicle (fibrofolliculomas). FLCN may be a tumor suppressor. May be involved in colorectal tumorigenesis. It may be involved in energy and/or nutrient sensing through the AMPK and mTOR signaling pathways.

FLCN Antibody (Center) - References

- Khoo,S.K., et.al., J. Med. Genet. 39 (12), 906-912 (2002)
Shin,J.H., et.al., J. Med. Genet. 40 (5), 364-367 (2003)