

DERA Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8764a**Specification**

DERA Antibody (N-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	O9Y315
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35231
Antigen Region	12-41

DERA Antibody (N-term) - Additional Information**Gene ID** 51071**Other Names**Deoxyribose-phosphate aldolase, DERA, 2-deoxy-D-ribose 5-phosphate aldolase,
Phosphodeoxyriboaldolase, Deoxyriboaldolase, DERA**Target/Specificity**

This DERA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 12-41 amino acids from the N-terminal region of human DERA.

DilutionWB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DERA Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DERA Antibody (N-term) - Protein Information**Name** DERA

Function Catalyzes a reversible aldol reaction between acetaldehyde and D-glyceraldehyde 3-phosphate to generate 2-deoxy-D-ribose 5-phosphate. Participates in stress granule (SG) assembly. May allow ATP production from extracellular deoxyinosine in conditions of energy deprivation.

Cellular Location

Cytoplasm. Cytoplasmic granule. Nucleus. Note=Recruited to stress granules but not to processing bodies upon arsenite or clotrimazole treatment or energy deprivation.

Tissue Location


Mainly expressed in liver, lung and colon.

DERA Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


DERA Antibody (N-term) - Images



Western blot analysis of DERA Antibody (N-term) (Cat. #AP8764a) in A375 cell line lysates (35ug/lane).DERA (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human skin reacted with DERA Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



DERA Antibody (N-term) (Cat. #AP8764a) flow cytometric analysis of A375 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

DERA Antibody (N-term) - References

Sgarrella, F., et al., Comp. Biochem. Physiol. B, Biochem. Mol. Biol. 117 (2), 253-257 (1997)