

**COQ3 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8765C****Specification**

---

**COQ3 Antibody (Center) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">O9NZJ6</a>
Other Accession	<a href="#">Q3T131</a>
Reactivity	Human
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	41054
Antigen Region	172-201

**COQ3 Antibody (Center) - Additional Information****Gene ID** 51805**Other Names**

Hexaprenyldihydroxybenzoate methyltransferase, mitochondrial, 2-polyprenyl-6-hydroxyphenol methylase, 4-dihydroxy-5-hexaprenylbenzoate methyltransferase, DHHB methyltransferase, DHHB-MT, DHHB-MTase, 3-demethylubiquinone-10 3-methyltransferase, Dihydroxyhexaprenylbenzoate methyltransferase, COQ3

**Target/Specificity**

This COQ3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 172-201 amino acids from the Central region of human COQ3.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

COQ3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**COQ3 Antibody (Center) - Protein Information**

**Name** COQ3 {ECO:0000255|HAMAP-Rule:MF\_03190}

**Function** O-methyltransferase that catalyzes the 2 O-methylation steps in the ubiquinone biosynthetic pathway.

**Cellular Location**


Mitochondrion inner membrane {ECO:0000255|HAMAP- Rule:MF\_03190}; Peripheral membrane protein {ECO:0000255|HAMAP- Rule:MF\_03190}; Matrix side {ECO:0000255|HAMAP-Rule:MF\_03190}

**COQ3 Antibody (Center) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**COQ3 Antibody (Center) - Images**



Western blot analysis of COQ3 Antibody (Center) (Cat. #AP8765c) in K562 cell line lysates (35ug/lane). COQ3 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human skeletal muscle reacted with COQ3 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



COQ3 Antibody (Center) (Cat.#AP8765c) FC analysis of ZR-75-1 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### COQ3 Antibody (Center) - Background

Ubiquinone, also known as coenzyme Q, or Q, is a critical component of the electron transport pathways of both eukaryotes and prokaryotes (Jonassen and Clarke, 2000 [PubMed 10777520]). This lipid consists of a hydrophobic isoprenoid tail and a quinone head group. The tail varies in length depending on the organism, but its purpose is to anchor coenzyme Q to the membrane. The quinone head group is responsible for the activity of coenzyme Q in the respiratory chain. COQ3 is an O-methyltransferase required for 2 steps in the biosynthetic pathway of coenzyme Q. This enzyme methylates an early coenzyme Q intermediate, 3,4-dihydroxy-5-polyprenylbenzoic acid, as well as the final intermediate in the pathway, converting demethyl-ubiquinone to coenzyme Q. The COQ3 is also capable of methylating the distinct prokaryotic early intermediate 2-hydroxy-6-polyprenyl phenol.

### COQ3 Antibody (Center) - References

Olsen,J.V., et.al., Cell 127 (3), 635-648 (2006)