10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CYP7B1 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8786c

Specification

CYP7B1 Antibody (Center) - Product Information

Application	**IF, WB, IHC-P, FC,E**
Primary Accession	O75881
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**58256**
Antigen Region	**252-281**

CYP7B1 Antibody (Center) - Additional Information

Gene ID 9420

Other Names
25-hydroxycholesterol 7-alpha-hydroxylase, Cytochrome P450 7B1, Oxysterol 7-alpha-hydroxylase, CYP7B1

Target/Specificity
This CYP7B1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 252-281 amino acids from the Central region of human CYP7B1.

Dilution
IF~~1:10~50
WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
CYP7B1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

CYP7B1 Antibody (Center) - Protein Information

Name CYP7B1 {ECO:0000303|PubMed:24491228, ECO:0000312|HGNC:HGNC:2652}

Function A cytochrome P450 monooxygenase involved in the metabolism of endogenous oxysterols and steroid hormones, including neurosteroids (PubMed:10588945, PubMed:24491228). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR; NADPH-ferrihemoprotein reductase) (PubMed:10588945, PubMed:24491228). Catalyzes the hydroxylation of carbon hydrogen bonds of steroids with a preference for 7-alpha position (PubMed:10588945, PubMed:24491228). Usually metabolizes steroids carrying a hydroxy group at position 3, functioning as a 3- hydroxy steroid 7-alpha hydroxylase (PubMed:24491228). Hydroxylates oxysterols, including 25-hydroxycholesterol and (25R)-cholest-5-ene- 3beta,26-diol toward 7-alpha hydroxy derivatives, which may be transported to the liver and converted to bile acids (PubMed:9802883, PubMed:10588945). Via its product 7-alpha,25-dihydroxycholesterol, a ligand for the chemotactic G protein-coupled receptor GPR183/EBI2, regulates B cell migration in germinal centers of lymphoid organs, thus guiding efficient maturation of plasma B cells and overall antigen- specific humoral immune response (By similarity). 7-alpha hydroxylates neurosteroids, including 3beta-hydroxyandrost-5-en-17-one (dehydroepiandrosterone) and pregnenolone, both involved in hippocampus-associated memory and learning (PubMed:24491228). Metabolizes androstanoids toward 6- or 7-alpha hydroxy derivatives (PubMed:24491228).

Cellular Location
Endoplasmic reticulum membrane; Multi-pass membrane protein. Microsome membrane; Multi-pass membrane protein

Tissue Location
Widely expressed. Expressed in brain, testis, ovary, prostate, liver, colon, kidney, small intestine, thymus and spleen.


CYP7B1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CYP7B1 Antibody (Center) - Images


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Confocal immunofluorescent analysis of CYP7B1 Antibody (Center) (Cat. #AP8786c) with Hela cell followed by Alexa Fluor® 488-conjugated goat anti-rabbit lgG (green). DAPI was used to stain the cell nuclear (blue).




Western blot analysis of CYP7B1 Antibody (Center) (Cat. #AP8786c) in Hela cell line lysates (35ug/lane). CYP7B1 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with CYP7B1 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



CYP7B1 Antibody (Center) (Cat. #AP8786c) flow cytometric analysis of Hela cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

CYP7B1 Antibody (Center) - Background

CYP7B1 is a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This endoplasmic reticulum membrane protein catalyzes the first reaction in the cholesterol catabolic pathway of extrahepatic tissues, which converts cholesterol to bile acids. This enzyme likely plays a minor role in total bile acid synthesis, but may also be involved in the development of atherosclerosis, neurosteroid metabolism and sex hormone synthesis.

CYP7B1 Antibody (Center) - References

Schwarz,M., et.al., Curr. Opin. Lipidol. 9 (2), 113-118 (1998)