

SOCS1 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8790A**Specification**

SOCS1 Antibody (N-term) - Product Information

Application	IF, WB, IHC-P, FC,E
Primary Accession	O15524
Reactivity	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	35-66

SOCS1 Antibody (N-term) - Additional Information**Gene ID** 8651**Other Names**

Suppressor of cytokine signaling 1, SOCS-1, JAK-binding protein, JAB, STAT-induced STAT inhibitor 1, SSI-1, Tec-interacting protein 3, TIP-3, SOCS1, SSI1, TIP3

Target/Specificity

This SOCS1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 35-66 amino acids from the N-terminal region of human SOCS1.

Dilution

IF~~1:10~50
WB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SOCS1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SOCS1 Antibody (N-term) - Protein Information**Name** SOCS1

Synonyms SSI1, TIP3 {ECO:0000303|PubMed:9341160}

Function Essential negative regulator of type I and type II interferon (IFN) signaling, as well as that of other cytokines, including IL2, IL4, IL6 and leukemia inhibitory factor (LIF) (PubMed:[32499645](#), PubMed:[33087723](#)). Downregulates cytokine signaling by inhibiting the JAK/STAT signaling pathway. Acts by binding to JAK proteins and to IFNGR1 and inhibiting their kinase activity. In vitro, suppresses Tec protein-tyrosine activity (PubMed:[9341160](#)). Regulates IFN-gamma (IFNG)- mediated sensory neuron survival (By similarity). Probable substrate recognition component of an ECS (Elongin BC-CUL2/5-SOCS-box protein) E3 ubiquitin ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:[11278610](#), PubMed:[11313480](#)).

Cellular Location

Nucleus. Cytoplasmic vesicle. Note=Detected in perinuclear cytoplasmic vesicles upon interaction with FGFR3

Tissue Location


Expressed in all tissues with high expression in spleen, small intestine and peripheral blood leukocytes

SOCS1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SOCS1 Antibody (N-term) - Images




Confocal immunofluorescent analysis of SOCS1 Antibody (N-term) (Cat. #AP8790a) with 293 cell followed by Alexa Fluor[®]488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the


cell nuclear (blue).




All lanes : Anti-SOCS1 Antibody (N-term) at 1:2000 dilution Lane 1: mouse thymus lysates Lane 2: rat spleen lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 24 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of SOCS1 Antibody (N-term) (Cat. #AP8790a) in mouse kidney tissue lysates (35ug/lane). SOCS1 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human colon carcinoma reacted with SOCS1 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



SOCS1 Antibody (N-term) (Cat. #AP8790a) flow cytometric analysis of WiDr cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

SOCS1 Antibody (N-term) - Background

SOCS1 is a member of the STAT-induced STAT inhibitor (SSI), also known as suppressor of cytokine signaling (SOCS), family. SSI family members are cytokine-inducible negative regulators of cytokine signaling.

SOCS1 Antibody (N-term) - References

Starr, R., et al., Nature 387 (6636), 917-921 (1997) Minamoto, S., et al., Biochem. Biophys. Res. Commun. 237 (1), 79-83 (1997)