

SELENBP1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP8869b

Specification

SELENBP1 Antibody (C-term) - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW Antigen Region WB, FC,E <u>Q13228</u> <u>NP_003935</u> Human, Mouse Rabbit Polyclonal Rabbit IgG 52391 317-344

SELENBP1 Antibody (C-term) - Additional Information

Gene ID 8991

Other Names Selenium-binding protein 1, 56 kDa selenium-binding protein, SBP56, SP56, SELENBP1, SBP

Target/Specificity

This SELENBP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 317-344 amino acids from the C-terminal region of human SELENBP1.

Dilution WB~~1:1000 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SELENBP1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SELENBP1 Antibody (C-term) - Protein Information

Name SELENBP1

Synonyms SBP



Function Catalyzes the oxidation of methanethiol, an organosulfur compound known to be produced in substantial amounts by gut bacteria (PubMed:<u>29255262</u>). Selenium-binding protein which may be involved in the sensing of reactive xenobiotics in the cytoplasm. May be involved in intra-Golgi protein transport (By similarity).

Cellular Location

Nucleus. Cytoplasm, cytosol Membrane {ECO:0000250|UniProtKB:Q8VIF7}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q8VIF7}. Note=May associate with Golgi membrane (By similarity). May associate with the membrane of autophagosomes (By similarity). {ECO:0000250|UniProtKB:Q8VIF7}

Tissue Location


Widely expressed. Highly expressed in liver, lung, colon, prostate, kidney and pancreas. In brain, present both in neurons and glia (at protein level). Down-regulated in lung adenocarcinoma, colorectal carcinoma and ovarian cancer. Two-fold up-regulated in brain and blood from schizophrenia patients.

SELENBP1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

SELENBP1 Antibody (C-term) - Images




Western blot analysis of SELENBP1 Antibody (C-term) (Cat. #AP8869b) in HL-60 cell line lysates (35ug/lane). SELENBP1 (arrow) was detected using the purified Pab.






Western blot analysis of SELENBP1 Antibody (C-term) (Cat. #AP8869b) in mouse lung tissue lysates (35ug/lane). SELENBP1 (arrow) was detected using the purified Pab.



Western blot analysis of SELENBP1 (arrow) using rabbit polyclonal SELENBP1 Antibody (C-term) (Cat. #AP8869b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the SELENBP1 gene.



SELENBP1 Antibody (C-term) (Cat. #AP8869b) flow cytometric analysis of HL-60 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

SELENBP1 Antibody (C-term) - Background

SELENBP1 is the selenium-binding protein family. Selenium is an essential nutrient that exhibits potent anticarcinogenic properties, and deficiency of selenium may cause certain neurologic



diseases. It has been proposed that the effects of selenium in preventing cancer and neurologic diseases may be mediated by selenium-binding proteins.

SELENBP1 Antibody (C-term) - References

Kanazawa, T., et.al., Schizophr. Res. 113 (2-3), 268-272 (2009)