

UNC13B Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8905c**Specification**

UNC13B Antibody (Center) - Product Information

Application	WB, FC,E
Primary Accession	O14795
Other Accession	Q9Z1N9
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	1062-1091

UNC13B Antibody (Center) - Additional Information**Gene ID** 10497**Other Names**

Protein unc-13 homolog B, Munc13-2, munc13, UNC13B, UNC13

Target/Specificity

This UNC13B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1062-1091 amino acids from the Central region of human UNC13B.

Dilution

WB~~1:1000

FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

UNC13B Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

UNC13B Antibody (Center) - Protein Information**Name** UNC13B ([HGNC:12566](#))**Synonyms** UNC13

Function Plays a role in vesicle maturation during exocytosis as a target of the diacylglycerol second messenger pathway. Is involved in neurotransmitter release by acting in synaptic vesicle priming prior to vesicle fusion and participates in the activity-dependent refilling of readily releasable vesicle pool (RRP) (By similarity). Essential for synaptic vesicle maturation in a subset of excitatory/glutamatergic but not inhibitory/GABA-mediated synapses (By similarity). In collaboration with UNC13A, facilitates neuronal dense core vesicles fusion as well as controls the location and efficiency of their synaptic release (By similarity).

Cellular Location

Cytoplasm. Membrane; Peripheral membrane protein. Cell membrane Synapse. Note=Localized to synapses. Translocated to the plasma membrane in response to phorbol ester binding (By similarity)

Tissue Location


Expressed in kidney cortical epithelial cells and brain.

UNC13B Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UNC13B Antibody (Center) - Images



Western blot analysis of UNC13B Antibody (Center) (Cat. #AP8905c) in 293 cell line lysates (35ug/lane). UNC13B (arrow) was detected using the purified Pab.



UNC13B Antibody (Center) (Cat. #AP8905c) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

UNC13B Antibody (Center) - Background

UNC13B plays a role in vesicle maturation during exocytosis as a target of the diacylglycerol second messenger pathway. It is involved in neurotransmitter release by acting in synaptic vesicle priming prior to vesicle fusion and participates in the activity-dependent refilling of readily releasable vesicle pool (RRP). It is essential for synaptic vesicle maturation in a subset of excitatory/glutamatergic but not inhibitory/GABA-mediated synapses (By similarity).

UNC13B Antibody (Center) - References

Sjoeblom T., et.al., Science 314:268-274(2006).
Yu L.-R., Zhu Z., et.al., J. Proteome Res. 6:4150-4162(2007).

UNC13B Antibody (Center) - Citations

- [Size-dependent mechanism of cargo sorting during lysosome-phagosome fusion is controlled by Rab34.](#)